

DUSP13-L184 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8455b

Product Information

WB, IHC-P, E
<u>Q9UII6</u>
Human
Rabbit
Polyclonal
Rabbit IgG
22149
169-198

Additional Information

Gene ID	51207
Other Names	Dual specificity protein phosphatase 13 isoform B, DUSP13B, Dual specificity phosphatase SKRP4, Testis- and skeletal-muscle-specific DSP, DUSP13, DUSP13B, TMDP
Target/Specificity	This DUSP13-L184 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 169-198 amino acids from the C-terminal region of human DUSP13-L184.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DUSP13-L184 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DUSP13B (<u>HGNC:19681</u>)
Synonyms	DUSP13, SKRP4, TMDP
Function	Dual specificity phosphatase that dephosphorylates MAPK8/JNK and MAPK14/p38, but not MAPK1/ERK2, in vitro (PubMed: <u>21360282</u>). Exhibits

	intrinsic phosphatase activity towards both phospho- seryl/threonyl and -tyrosyl residues, with similar specific activities in vitro (PubMed: <u>10585869</u>).
Tissue Location	Highly expressed in the testis (at protein level) (PubMed:10585869, PubMed:15252030). Also found in the skeletal muscle (PubMed:15252030).

Background

Dual-specificity phosphatases, a subfamily of protein-tyrosine phosphatases, play important roles in signal transduction, cell cycle progression, and tumor suppression. The cDNA encoding a novel phosphatase, PIR1, phosphatase that interacts with RNA/RNP complex 1. Sequence analysis revealed that the predicted 329-amino acid protein has homology to several dual-specificity phosphatases and contains 2 stretches of arginine-rich sequence similar to those found in some RNA-binding proteins. In vitro, recombinant protein displays protein-tyrosine phosphatase activity and binds directly to RNA.

References

Nakamura K., Shima H., Watanabe M., Haneji T., Kikuchi K.Biochem. J. 344:819-825(1999). Deloukas et al. Nature 429:375-381(2004). Strausberg et al. Proc. Natl. Acad. Sci. U.S.A. 99:16899-16903(2002).

Images



Citations

• Regulation of MITF stability by the USP13 deubiquitinase.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.