

PP5 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8476b

Product Information

Application	WB, IHC-P, E
Primary Accession	P53041
Other Accession	P53042 , Q60676 , NP_006238
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB5864
Calculated MW	56879
Antigen Region	445-475

Additional Information

Gene ID	5536
Other Names	Serine/threonine-protein phosphatase 5, PP5, Protein phosphatase T, PP-T, PPT, PPP5C, PPP5
Target/Specificity	This PP5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 445-475 amino acids from the C-terminal region of human PP5 .
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PP5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPP5C
Synonyms	PPP5

Function	Serine/threonine-protein phosphatase that dephosphorylates a myriad of proteins involved in different signaling pathways including the kinases CSNK1E, ASK1/MAP3K5, PRKDC and RAF1, the nuclear receptors NR3C1, PPARG, ESR1 and ESR2, SMAD proteins and TAU/MAPT (PubMed:14734805, PubMed:14764652, PubMed:14871926, PubMed:15383005, PubMed:15546861, PubMed:16260606, PubMed:16790549, PubMed:16892053, PubMed:19176521, PubMed:19948726, PubMed:21144835, PubMed:22399290, PubMed:22781750, PubMed:23102700, PubMed:30699359, PubMed:9000529). Implicated in wide ranging cellular processes, including apoptosis, differentiation, DNA damage response, cell survival, regulation of ion channels or circadian rhythms, in response to steroid and thyroid hormones, calcium, fatty acids, TGF-beta as well as oxidative and genotoxic stresses (PubMed:14734805, PubMed:14764652, PubMed:14871926, PubMed:15383005, PubMed:15546861, PubMed:16260606, PubMed:16790549, PubMed:16892053, PubMed:19176521, PubMed:19948726, PubMed:21144835, PubMed:22399290, PubMed:22781750, PubMed:23102700, PubMed:30699359, PubMed:9000529). Participates in the control of DNA damage response mechanisms such as checkpoint activation and DNA damage repair through, for instance, the regulation ATM/ATR-signaling and dephosphorylation of PRKDC and TP53BP1 (PubMed:14871926, PubMed:16260606, PubMed:21144835). Inhibits ASK1/MAP3K5-mediated apoptosis induced by oxidative stress (PubMed:23102700). Plays a positive role in adipogenesis, mainly through the dephosphorylation and activation of PPARG transactivation function (By similarity). Also dephosphorylates and inhibits the anti- adipogenic effect of NR3C1 (By similarity). Regulates the circadian rhythms, through the dephosphorylation and activation of CSNK1E (PubMed:16790549). May modulate TGF-beta signaling pathway by the regulation of SMAD3 phosphorylation and protein expression levels (PubMed:22781750). Dephosphorylates and may play a role in the regulation of TAU/MAPT (PubMed:15546861). Through their dephosphorylation, may play a role in the regulation of ions channels such as KCNH2 (By similarity). Dephosphorylate FNIP1, disrupting interaction with HSP90AA1/Hsp90 (PubMed:30699359).
Cellular Location	Nucleus. Cytoplasm. Cell membrane. Note=Predominantly nuclear (PubMed:15383005). But also present in the cytoplasm (PubMed:15383005) Translocates from the cytoplasm to the plasma membrane in a RAC1-dependent manner (PubMed:19948726).
Tissue Location	Ubiquitous..


Background

PPP5C belongs to the PPP phosphatase family, PP-T subfamily, and may play a role in the regulation of RNA biogenesis and/or mitosis. In vitro,PPP5C dephosphorylates serine residues of skeletal muscle phosphorylase and histone H1.


References

- Yang, J., et al., EMBO J. 24(1):1-10 (2005).
 Zhou, G., et al., J. Biol. Chem. 279(45):46595-46605 (2004).
 Huang, S., et al., J. Biol. Chem. 279(35):36490-36496 (2004).
 Swingle, M.R., et al., J. Biol. Chem. 279(32):33992-33999 (2004).
 Wechsler, T., et al., Proc. Natl. Acad. Sci. U.S.A. 101(5):1247-1252 (2004).


Images



Western blot analysis of PP5(Cat.#AP8476b) in A375, A2058 cell line lysates (35ug/lane). PP5 (arrow) was detected using the purified Pab.



All lanes : Anti-PP5 Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.