

# IGFBP2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8588B

## Product Information

---

<b>Application</b>	IHC-P, IF, FC, WB, E
<b>Primary Accession</b>	<a href="#">P18065</a>
<b>Other Accession</b>	<a href="#">P24853</a> , <a href="#">P13384</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Predicted</b>	Bovine, Pig
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB22493
<b>Calculated MW</b>	34814
<b>Antigen Region</b>	277-305

## Additional Information

---

<b>Gene ID</b>	3485
<b>Other Names</b>	Insulin-like growth factor-binding protein 2, IBP-2, IGF-binding protein 2, IGFBP-2, IGFBP2, BP2, IBP2
<b>Target/Specificity</b>	This IGFBP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 277-305 amino acids from the C-terminal region of human IGFBP2.
<b>Dilution</b>	IHC-P~~1:100~500 IF~~1:10~50 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	IGFBP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	IGFBP2
<b>Synonyms</b>	BP2, IBP2

<b>Function</b>	Multifunctional protein that plays a critical role in regulating the availability of IGFs such as IGF1 and IGF2 to their receptors and thereby regulates IGF-mediated cellular processes including proliferation, differentiation, and apoptosis in a cell-type specific manner (PubMed: <a href="#">18563800</a> , PubMed: <a href="#">38796567</a> ). Functions coordinately with receptor protein tyrosine phosphatase beta/PTPRB and the IGF1 receptor to regulate IGF1-mediated signaling by stimulating the phosphorylation of PTEN leading to its inactivation and AKT1 activation (PubMed: <a href="#">22869525</a> ). Plays a positive role in cell migration via interaction with integrin alpha5/ITGA5 through an RGD motif (PubMed: <a href="#">16569642</a> ). Additionally, interaction with ITGA5/ITGB1 enhances the adhesion of endothelial progenitor cells to endothelial cells (PubMed: <a href="#">26076738</a> ). Upon mitochondrial damage, facilitates apoptosis with ITGA5 of podocytes, and then activates the phosphorylation of focal adhesion kinase (FAK)-mediated mitochondrial injury (PubMed: <a href="#">38796567</a> ).
<b>Cellular Location</b>	Secreted


## Background

IGFBP2-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.

## References

Arafat,A.M.,et.al.,J. Clin. Endocrinol. Metab. 94 (12), 5093-5101 (2009)  
Yazawa,T.,et.al.,Am. J. Pathol. 175 (3), 976-987 (2009)

## Images



Anti-IGFBP2 Antibody (C-term) at 1:1000 dilution + T47D whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.