

# NQO2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8618c

#### **Product Information**

**Application** WB, E **Primary Accession** P16083

**Reactivity** Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB21132
Calculated MW 25919
Antigen Region 141-168

### **Additional Information**

**Gene ID** 4835

Other Names Ribosyldihydronicotinamide dehydrogenase [quinone], NRH dehydrogenase

[quinone] 2, NRH:quinone oxidoreductase 2, Quinone reductase 2, QR2,

NQO2, NMOR2

**Target/Specificity** This NQO2 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 141-168 amino acids from the Central

region of human NQO2.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** NQO2 Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name NQO2

Synonyms NMOR2

**Function** The enzyme apparently serves as a quinone reductase in connection with

conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.

**Cellular Location** 

Cytoplasm.


# **Background**

NQO2 apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.

## References

Foster, C.E., et.al., Biochemistry 38 (31), 9881-9886 (1999) Wu, K., Knox, R., et.al., Arch. Biochem. Biophys. 347 (2), 221-228 (1997)

## **Images**



Western blot analysis of NQO2 Antibody (Center) (Cat. #AP8618c) in mouse liver tissue lysates (35ug/lane). NQO2 (arrow) was detected using the purified Pab.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.