

# GSTA1 Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8697a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P08263</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB22670
<b>Calculated MW</b>	25631

## Additional Information

---

<b>Gene ID</b>	2938
<b>Other Names</b>	Glutathione S-transferase A1, GST HA subunit 1, GST class-alpha member 1, GST-epsilon, GSTA1-1, GTH1, Glutathione S-transferase A1, N-terminally processed, GSTA1
<b>Target/Specificity</b>	This GSTA1 antibody is generated from rabbits immunized with GSTA1 recombinant protein.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	GSTA1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GSTA1
<b>Function</b>	Glutathione S-transferase that catalyzes the nucleophilic attack of the sulfur atom of glutathione on the electrophilic groups of a wide range of exogenous and endogenous compounds (Probable). Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed: <a href="#">9084911</a> ). It also catalyzes the isomerization of

D5-androstene-3,17-dione (AD) into D4-androstene- 3,17-dione and may therefore play an important role in hormone biosynthesis (PubMed:[11152686](#)). Through its glutathione-dependent peroxidase activity toward the fatty acid hydroperoxide (13S)-hydroperoxy-(9Z,11E)-octadecadienoate/13-HPODE it is also involved in the metabolism of oxidized linoleic acid (PubMed:[16624487](#)).

**Cellular Location** Cytoplasm.

**Tissue Location** Liver.


## Background

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes function in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione.

## References

Bogaards,J.J., et.al., Biochem. J. 286 (PT 2), 383-388 (1992)  
Mikhal'chik,E.V., et.al., Bull. Exp. Biol. Med. 147 (6), 753-756 (2009)

## Images



All lanes : Anti-GSTA1 Antibody at 1:4000 dilution Lane 1: HepG2 whole cell lysate Lane 2: Caco2 whole cell lysate Lane 3: LNCaP whole cell lysate Lane 4: PANC-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.