

# FPRL2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8793c

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P25089</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB21312
<b>Calculated MW</b>	39965
<b>Antigen Region</b>	307-333

## Additional Information

---

<b>Gene ID</b>	2359
<b>Other Names</b>	N-formyl peptide receptor 3, FMLP-related receptor II, FMLP-R-II, Formyl peptide receptor-like 2, FPR3, FPRH1, FPRL2
<b>Target/Specificity</b>	This FPRL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 307-333 amino acids from the Central region of human FPRL2.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	FPRL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	FPR3
<b>Synonyms</b>	FPRH1, FPRL2
<b>Function</b>	Low affinity receptor for N-formyl-methionyl peptides, which are powerful

neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. Acts as a receptor for humanin (PubMed:[15465011](#)).

**Cellular Location** Cell membrane; Multi-pass membrane protein.

**Tissue Location** Detected in various tissues with highest expression in lung.

## Background

---

Low affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.


## References

---


Yang,D., et.al., J. Leukoc. Biol. 72 (3), 598-607 (2002)

## Images

---




Western blot analysis of FPRL2 Antibody (Center) (Cat. #AP8793c) in NCI-H460 cell line lysates (35ug/lane). FPRL2 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with FPRL2 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

FPRL2 Antibody (Center) (Cat. #AP8793c) flow cytometry analysis of NCI-H460 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.