

GAL3ST1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8884c

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q99999
Reactivity	Human, Mouse
Predicted	Rat, Bovine, Canine, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB21441
Calculated MW	48764
Antigen Region	88-116

Additional Information

Gene ID	9514
Other Names	Galactosylceramide sulfotransferase, GalCer sulfotransferase, 3'-phosphoadenosine-5'-phosphosulfate:GalCer sulfotransferase, 3'-phosphoadenylylsulfate:galactosylceramide 3'-sulfotransferase, Cerebroside sulfotransferase, GAL3ST1, CST
Target/Specificity	This GAL3ST1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-116 amino acids from the Central region of human GAL3ST1.
Dilution	WB~~1:2000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GAL3ST1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GAL3ST1 (HGNC:24240)
-------------	--

Function	Catalyzes the transfer of a sulfate group to position 3 of non-reducing beta-galactosyl residues in glycerolipids and sphingolipids, therefore participates in the biosynthesis of sulfoglycolipids (PubMed: 8830034 , PubMed: 9030544). Catalyzes the synthesis of galactosylceramide sulfate (sulfatide), a major lipid component of the myelin sheath and of monogalactosylalkylacylglycerol sulfate (seminolipid), present in spermatocytes (PubMed: 8830034). Seems to prefer beta-glycosides at the non-reducing termini of sugar chains attached to a lipid moiety (PubMed: 8830034). Also acts on lactosylceramide, galactosyl 1-alkyl-2-sn-glycerol and galactosyl diacylglycerol (in vitro) (PubMed: 8830034).
Cellular Location	Golgi apparatus membrane; Single- pass type II membrane protein
Tissue Location	Expressed in kidney proximal tubule, gastric mucosa and adenocarcinoma (PubMed:10785389, PubMed:9030544). Highly expressed in renal cell carcinoma cell lines (PubMed:8830034, PubMed:9030544)


Background

Sulfonation, an important step in the metabolism of many drugs, xenobiotics, hormones, and neurotransmitters, is catalyzed by sulfotransferases. GAL3ST1 is galactosylceramide sulfotransferase which catalyzes the conversion between 3'-phosphoadenylylsulfate + a galactosylceramide to adenosine 3',5'-bisphosphate + galactosylceramide sulfate. Activity of this sulfotransferase is enhanced in renal cell carcinoma.


References

Siegrist,H.P., et.al., Biochim. Biophys. Acta 489 (1), 58-63 (1977)
Stein,C., et.al., J. Biol. Chem. 264 (2), 1252-1259 (1989)


Images



Western blot analysis of GAL3ST1 Antibody (Center) (Cat. #AP8884c) in mouse cerebellum tissue lysates (35ug/lane). GAL3ST1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with GAL3ST1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



GAL3ST1 Antibody (Center) (Cat. #AP8884c) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.