

# Lamin B1 Antibody

Rabbit mAb

Catalog # AP90034

## Product Information

<b>Application</b>	WB, IHC, IF, ICC, IP, IHF
<b>Primary Accession</b>	<a href="#">P20700</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	LMN; ADLD; LMN2; LMNB; Lamin B1;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	66408


## Additional Information

<b>Dilution</b>	WB 1:3000~1:10000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human Lamin B1
<b>Description</b>	The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


## Protein Information

<b>Name</b>	LMNB1
<b>Synonyms</b>	LMN2, LMNB
<b>Function</b>	Lamins are intermediate filament proteins that assemble into a filamentous meshwork, and which constitute the major components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane (PubMed: <a href="#">28716252</a> , PubMed: <a href="#">32910914</a> ). Lamins provide a framework for the nuclear envelope, bridging the nuclear envelope and chromatin, thereby playing an important role in nuclear assembly, chromatin organization, nuclear membrane and telomere dynamics (PubMed: <a href="#">28716252</a> , PubMed: <a href="#">32910914</a> ). The structural integrity of the lamina is strictly controlled by the cell cycle, as seen by the disintegration and formation of the nuclear envelope in prophase and telophase, respectively (PubMed: <a href="#">28716252</a> , PubMed: <a href="#">32910914</a> ).


Images



All lanes use AP90034 Lamin B1 Antibody at 1:10000 dilution for 1 hour at room temperature.



All lanes use AP90034 Lamin B1 Antibody at 1:20000 dilution for 1 hour at room temperature.



All lanes use AP90034 Lamin B1 Antibody at 1:10000 dilution for 1 hour at room temperature.

Image not found : 202311/AP90034-IHC.jpg

Immunohistochemical analysis of paraffin-embedded rat liver, using Lamin B1 Antibody.

Image not found : 202311/AP90034-IF.jpg

Immunofluorescent analysis of HepG2 cells, using Lamin B1 Antibody.

Image not found : 202311/AP90034-wb6.jpg

Zinc inhibited LPS-induced inflammatory responses by upregulating A20 expression in microglia BV2 cells.  
-Journal of Affective Disorders

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.