10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Cubilin Antibody

Rabbit mAb
Catalog # AP90069

Product Information

Application	WB
Primary Accession	O60494
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Cubilin; Cubn; IFCR; MGA1; Cubilin precursor; megaloblastic anemia 1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	398736

Additional Information


Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Cubilin
Description	Cotransporter which plays a role in lipoprotein, vitamin and iron metabolism, by facilitating their uptake. Binds to ALB, MB, Kappa and lambda-light chains, TF, hemoglobin, GC, SCGB1A1, APOA1, high density lipoprotein, and the GIF-cobalamin complex. The binding of all ligands requires calcium. Serves as important transporter in several absorptive epithelia, including intestine, renal proximal tubules and embryonic yolk sac.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	CUBN
Synonyms	IFCR
Function	Endocytic receptor which plays a role in lipoprotein, vitamin and iron metabolism by facilitating their uptake (PubMed:10371504, PubMed:11606717, PubMed:11717447, PubMed:14576052, PubMed:9572993). Acts together with LRP2 to mediate endocytosis of high-density lipoproteins, GC, hemoglobin, ALB, TF and SCGB1A1. Acts together with AMN to mediate endocytosis of the CBLIF-cobalamin complex (PubMed:14576052, PubMed:9572993). Binds to ALB, MB, Kappa and lambda-light chains, TF, hemoglobin, GC, SCGB1A1, APOA1, high density lipoprotein, and the CBLIF-cobalamin complex. Ligand binding requires calcium (PubMed:9572993). Serves as important transporter in several absorptive epithelia, including intestine, renal proximal tubules and embryonic yolk sac.

	May play an important role in the development of the peri-implantation embryo through internalization of APOA1 and cholesterol. Binds to LGALS3 at the maternal-fetal interface.
Cellular Location	Apical cell membrane {ECO:0000250|UniProtKB:Q9JLB4}; Peripheral membrane protein. Cell membrane; Peripheral membrane protein {ECO:0000305, ECO:0000305|PubMed:30523278}. Membrane, coated pit. Endosome. Lysosome membrane {ECO:0000250|UniProtKB:O70244}; Peripheral membrane protein. Note=Lacks a transmembrane domain and depends on interaction with AMN for location at the plasma membrane (PubMed:29402915, PubMed:30523278). Colocalizes with AMN and LRP2 in the endocytotic apparatus of epithelial cells (By similarity) {ECO:0000250|UniProtKB:O70244, ECO:0000269|PubMed:29402915, ECO:0000269|PubMed:30523278}
Tissue Location	Detected in kidney cortex (at protein level) (PubMed:9572993). Expressed in kidney proximal tubule cells, placenta, visceral yolk-sac cells and in absorptive intestinal cells. Expressed in the epithelium of intestine and kidney

Images



Western blot analysis of Cubilin expression in Human fetal kidney lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.