

# EXOC5 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9006B

#### **Product Information**

**Application** IHC-P, FC, WB, E

**Primary Accession** <u>000471</u> Other Accession Q8TDX5 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB22622 **Calculated MW** 81853 40-66 **Antigen Region** 

## **Additional Information**

**Gene ID** 10640

Other Names Exocyst complex component 5, Exocyst complex component Sec10, hSec10,

EXOC5, SEC10, SEC10L1

Target/Specificity This EXOC5 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 40-66 amino acids from the C-terminal

region of human EXOC5.

**Dilution** IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent

concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** EXOC5 Antibody (C-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name EXOC5

Synonyms SEC10, SEC10L1

**Function** Component of the exocyst complex involved in the docking of exocytic

vesicles with fusion sites on the plasma membrane.

**Cellular Location** Cytoplasm. Midbody. Note=Localization at the midbody requires the presence

of RALA, EXOC2 and EXOC3

Tissue Location Ubiquitous...


# **Background**

EXOC5 is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity.


## References

Kee,Y., et.al., Proc. Natl. Acad. Sci. U.S.A. 94 (26), 14438-14443 (1997) Hsu,S.C., et.al., Neuron 20 (6), 1111-1122 (1998)

# **Images**




Anti-EXOC5 Antibody (C-term) at 1:1000 dilution + NCI-H460 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 82 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with EXOC5 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

EXOC5 Antibody (C-term) (Cat. #AP9006b) flow cytometric analysis of NCI-H460 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.