

Topoisomerase II alpha Antibody

Rabbit mAb Catalog # AP90075

Product Information


Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IP <u>P11388</u> Rat, Human, Mouse Monoclonal DNA topoisomerase II; DNA topoisomerase II-alpha; TOP2; TOP2A; TP2A; alpha isozyme;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	174385

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human Topoisomerase II alpha DNA topoisomerase 2-alpha; DNA topoisomerase II, 170 kD; DNA topoisomerase II, alpha isozyme; TOP2; TOP2A; topoisomerase (DNA) II alpha 170kDa; TP2A.
Storage Condition and Buffer	

Protein Information

Name	ΤΟΡ2Α
Synonyms	TOP2
Function	Key decatenating enzyme that alters DNA topology by binding to two double-stranded DNA molecules, generating a double-stranded break in one of the strands, passing the intact strand through the broken strand, and religating the broken strand (PubMed: <u>17567603</u> , PubMed: <u>18790802</u> , PubMed: <u>22013166</u> , PubMed: <u>22323612</u>). May play a role in regulating the period length of BMAL1 transcriptional oscillation (By similarity).
Cellular Location	Cytoplasm. Nucleus, nucleoplasm. Nucleus. Nucleus, nucleolus
Tissue Location	Expressed in the tonsil, spleen, lymph node, thymus, skin, pancreas, testis, colon, kidney, liver, brain and lung (PubMed:9155056). Also found in high-grade lymphomas, squamous cell lung tumors and seminomas (PubMed:9155056)



Western blot analysis of Topoisomerase II alpha expression in (1) HeLa cell lysate; (2) Jurkat cell lysate.

Image not found : 202311/AP90075-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human colon cancer, using Topoisomerase II alpha Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.