

# active Caspase-3 Antibody

Rabbit mAb  
Catalog # AP90098

## Product Information

---

<b>Application</b>	WB, IHC, IF, ICC, IHF
<b>Primary Accession</b>	<a href="#">P42574</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	Apopain precursor; Cysteine protease CPP32; ICE3; CASP-3; CPP32; Caspase-3; SCA-1;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	31608

## Additional Information

---

<b>Dilution</b>	WB 1:500~1:2000 IHC 1:50~1:100 ICC/IF 1:50~1:100
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human active Caspase-3
<b>Description</b>	Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

---

<b>Name</b>	CASP3
<b>Synonyms</b>	CPP32 {ECO:0000303   PubMed:7983002}
<b>Function</b>	Thiol protease that acts as a major effector caspase involved in the execution phase of apoptosis (PubMed: <a href="#">18723680</a> , PubMed: <a href="#">20566630</a> , PubMed: <a href="#">23650375</a> , PubMed: <a href="#">35338844</a> , PubMed: <a href="#">35446120</a> , PubMed: <a href="#">7596430</a> ). Following cleavage and activation by initiator caspases (CASP8, CASP9 and/or CASP10), mediates execution of apoptosis by catalyzing cleavage of many proteins (PubMed: <a href="#">18723680</a> , PubMed: <a href="#">20566630</a> , PubMed: <a href="#">23650375</a> , PubMed: <a href="#">7596430</a> ). At the onset of apoptosis, it proteolytically cleaves poly(ADP-ribose) polymerase PARP1 at a '216-Asp- -Gly-217' bond (PubMed: <a href="#">10497198</a> , PubMed: <a href="#">16374543</a> , PubMed: <a href="#">7596430</a> , PubMed: <a href="#">7774019</a> ). Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain (By similarity). Cleaves

and activates caspase-6, -7 and -9 (CASP6, CASP7 and CASP9, respectively) (PubMed:[7596430](#)). Cleaves and inactivates interleukin-18 (IL18) (PubMed:[37993714](#), PubMed:[9334240](#)). Involved in the cleavage of huntingtin (PubMed:[8696339](#)). Triggers cell adhesion in sympathetic neurons through RET cleavage (PubMed:[21357690](#)). Cleaves DSG2 in response to apoptosis resulting in a loss of full length DSG2 at desmosome cell junctions and subsequent loss of cell-cell adhesion (PubMed:[17559062](#)). Also cleaves JUP in response to apoptosis (PubMed:[17559062](#)). Cleaves and inhibits serine/threonine-protein kinase AKT1 in response to oxidative stress (PubMed:[23152800](#)). Acts as an inhibitor of type I interferon production during virus-induced apoptosis by mediating cleavage of antiviral proteins CGAS, IRF3 and MAVS, thereby preventing cytokine overproduction (PubMed:[30878284](#)). Also involved in pyroptosis by mediating cleavage and activation of gasdermin-E (GSDME) (PubMed:[35338844](#), PubMed:[35446120](#)). Cleaves XRCC4 and phospholipid scramblase proteins XKR4, XKR8 and XKR9, leading to promote phosphatidylserine exposure on apoptotic cell surface (PubMed:[23845944](#), PubMed:[33725486](#)). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed:[36758104](#), PubMed:[36758106](#)).

#### Cellular Location


Cytoplasm.

#### Tissue Location

Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.

## Images

---



Western blot analysis of active Caspase-3 expression in Jurkat cell lysate treated with Camptothecin.

Image not found : 202311/AP90098-IF.jpg

Immunofluorescent analysis of HepG2 cells, using active Caspase-3 Antibody .

Image not found : 202311/AP90098-wb6.jpg

Combination of novel DR5 targeting agonistic scFv antibody TR2-3 with cisplatin shows enhanced synergistic antitumor activity in vitro and in vivo. -Biomedicine & Pharmacotherapy

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.