

# Phospho-c-Myc (T58 + S62) Antibody

Rabbit mAb Catalog # AP90113

#### **Product Information**

**Application** WB, IF, FC, ICC, IP

Primary Accession
Reactivity
Clonality
P01106
Rat, Human
Monoclonal

Other Names Myc proto-oncogene protein; c-myc;

IsotypeRabbit IgGHostRabbitCalculated MW50565

### **Additional Information**

**Dilution** WB 1:500~1:2000 ICC/IF 1:100~1:200 IP 1:50

**Purification** Affinity-chromatography

**Immunogen**A synthesized peptide derived from human Phospho-c-Myc (T58 + S62) **Description**Myc a proto-oncogenic transcription factor that plays a role in cell

proliferation, apoptosis and in the development of human tumors.. Seems to

activate the transcription of growth-related genes.

**Storage Condition and Buffer** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name MYC

Synonyms BHLHE39

**Function** Transcription factor that binds DNA in a non-specific manner, yet also

specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed: 24940000, PubMed: 25956029). Activates the transcription of growth-related genes (PubMed: 24940000, PubMed: 25956029). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis

(PubMed:<u>24940000</u>, PubMed:<u>25956029</u>). Regulator of somatic

reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA


polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production

of the PKM M2 isoform (PubMed: 20010808).

#### **Cellular Location**

Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Chromosome. Note=Association with chromatin is reduced by hyperphosphorylation (PubMed:30158517) Localization to the nucleolus is dependent on HEATR1 (PubMed:38225354)

## **Images**



Western blot analysis of Phospho-c-Myc (T58 + S62) in HEK293 cell lysate treated with EGF.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.