

Phospho-PP2A alpha (Y307) Antibody

Rabbit mAb Catalog # AP90124

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IF, ICC, IP, IHF <u>P67775</u> Rat, Human, Mouse Monoclonal eplication protein C; PP2A-alpha; Replication protein C; RP-C; PPP2CA; MGC786
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	35594

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human Phospho-PP2A alpha (Y307) PP2A is the major phosphatase for microtubule-associated proteins (MAPs). PP2A can modulate the activity of phosphorylase B kinase casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase. Cooperates with SGOL2 to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I (By similarity). Can dephosphorylate SV40 large T
Storage Condition and Buffer	antigen and p53/TP53. Activates RAF1 by dephosphorylating it at 'Ser-259'. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	PPP2CA
Function	Catalytic subunit of protein phosphatase 2A (PP2A), a serine/threonine phosphatase involved in the regulation of a wide variety of enzymes, signal transduction pathways, and cellular events (PubMed: <u>10801873</u> , PubMed: <u>12473674</u> , PubMed: <u>17245430</u> , PubMed: <u>22613722</u> , PubMed: <u>33243860</u> , PubMed: <u>34004147</u> , PubMed: <u>9920888</u>). PP2A is the major phosphatase for microtubule-associated proteins (MAPs) (PubMed: <u>22613722</u>). PP2A can modulate the activity of phosphorylase B kinase casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase (PubMed: <u>22613722</u>). Cooperates with SGO2 to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I (By similarity). Can dephosphorylate various proteins, such as SV40 large T antigen, AXIN1, p53/TP53, PIM3, WEE1 (PubMed: <u>10801873</u> ,

PubMed:<u>12473674</u>, PubMed:<u>17245430</u>, PubMed:<u>9920888</u>). Activates RAF1 by dephosphorylating it at 'Ser-259' (PubMed: 10801873). Mediates dephosphorylation of WEE1, preventing its ubiquitin-mediated proteolysis, increasing WEE1 protein levels, and promoting the G2/M checkpoint (PubMed:<u>33108758</u>). Mediates dephosphorylation of MYC; promoting its ubiguitin-mediated proteolysis: interaction with AMBRA1 enhances interaction between PPP2CA and MYC (PubMed:25438055). Mediates dephosphorylation of FOXO3; promoting its stabilization: interaction with AMBRA1 enhances interaction between PPP2CA and FOXO3 (PubMed:<u>30513302</u>). Catalyzes dephosphorylation of the pyrin domain of NLRP3, promoting assembly of the NLRP3 inflammasome (By similarity). Together with RACK1 adapter, mediates dephosphorylation of AKT1 at 'Ser-473', preventing AKT1 activation and AKT-mTOR signaling pathway (By similarity). Dephosphorylation of AKT1 is essential for regulatory T-cells (Treg) homeostasis and stability (By similarity). Catalyzes dephosphorylation of PIM3, promotinh PIM3 ubiquitination and proteasomal degradation (PubMed:12473674). Part of the striatin- interacting phosphatase and kinase (STRIPAK) complexes (PubMed:<u>33633399</u>). STRIPAK complexes have critical roles in protein (de)phosphorylation and are regulators of multiple signaling pathways including Hippo, MAPK, nuclear receptor and cytoskeleton remodeling (PubMed:33633399). Different types of STRIPAK complexes are involved in a variety of biological processes such as cell growth, differentiation, apoptosis, metabolism and immune regulation (PubMed:<u>33633399</u>). Key mediator of a quality checkpoint during transcription elongation as part of the Integrator-PP2A (INTAC) complex (PubMed:<u>33243860</u>, PubMed:<u>34004147</u>, PubMed:<u>37080207</u>). The INTAC complex drives premature transcription termination of transcripts that are unfavorably configured for transcriptional elongation: within the INTAC complex, PPP2CA catalyzes dephosphorylation of the C-terminal domain (CTD) of Pol II subunit POLR2A/RPB1 and SUPT5H/SPT5, thereby preventing transcriptional elongation (PubMed:<u>33243860</u>, PubMed:<u>34004147</u>, PubMed:37080207).

Cellular Location Cytoplasm. Nucleus. Chromosome. Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=In prometaphase cells, but not in anaphase cells, localizes at centromeres (PubMed:16541025). During mitosis, also found at spindle poles (PubMed:16541025). Centromeric localization requires the presence of SGO2 (By similarity). Recruited to chromatin and transcription pause-release checkpoint via its association with the Integrator complex (PubMed:33243860, PubMed:34004147). {ECO:0000250|UniProtKB:P63330, ECO:0000269|PubMed:16541025, ECO:0000269|PubMed:34004147}

Images



Western blot analysis of Phospho-PP2A alpha (Y307) in (1) Rat kidney lysate; (2) A431 cell lysate treated with EGF. Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.