

ENO1 Antibody

Rabbit mAb

Catalog # AP90166

Product Information

Application	WB, IF, ICC, IP
Primary Accession	P06733
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	ENO1;ENO1L1;MBP-1;MPB1;NNE;PPH;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	47169

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ENO1
Description	Enolase is an important glycolytic enzyme involved in the interconversion of 2-phosphoglycerate to phosphoenolpyruvate. Mammalian enolase exists as three subunits: enolase-1 (α -enolase), enolase-2 (γ -enolase) and enolase-3 (β -enolase) that can form both homo- and heterodimers. Expression of the enolase isoforms differs in a tissue specific manner.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	ENO1 (HGNC:3350)
Synonyms	ENO1L1, MBPB1, MPB1
Function	Enolase that catalyzes the conversion of 2-phosphoglycerate to phosphoenolpyruvate in glycolysis and the reverse reaction in gluconeogenesis (PubMed: 1369209 , PubMed: 29775581). Also involved in various processes such as growth control, hypoxia tolerance and allergic responses (PubMed: 10802057 , PubMed: 12666133 , PubMed: 2005901 , PubMed: 29775581). May also function in the intravascular and pericellular fibrinolytic system due to its ability to serve as a receptor and activator of plasminogen on the cell surface of several cell-types such as leukocytes and neurons (PubMed: 12666133). Stimulates immunoglobulin production (PubMed: 1369209).
Cellular Location	Cytoplasm. Cell membrane. Cytoplasm, myofibril, sarcomere, M line.

Note=Can translocate to the plasma membrane in either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form. ENO1 is localized to the M line

Tissue Location

The alpha/alpha homodimer is expressed in embryo and in most adult tissues. The alpha/beta heterodimer and the beta/beta homodimer are found in striated muscle, and the alpha/gamma heterodimer and the gamma/gamma homodimer in neurons

Images



Western blot analysis of ENO1 in (1)MCF-7 whole cell lysate; (2)Rat brain tissue lysate.

Image not found : 202311/AP90166-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human colon, using ENO1 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.