

Histone H3 (di methyl K4) Antibody

Rabbit mAb Catalog # AP90181

Product Information

Application WB, IHC, IF, FC, ICC, IP, IHF

Primary Accession P68431

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names Histone H3.1; Histone H3; HIST1H3A; H3K4me2;

IsotypeRabbit IgGHostRabbitCalculated MW15404

Additional Information

Dilution WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:50

Purification Affinity-chromatography

ImmunogenA synthesized peptide derived from human Methyl-Histone H3 (di K4)DescriptionH3 Core component of nucleosome. Nucleosomes wrap and compact DNA

into chromatin, limiting DNA accessibility to the cellular machineries which

require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal

stability.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name H3C1 (<u>HGNC:4766</u>)

Synonyms H3FA, HIST1H3A


Function Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

Cellular Location Nucleus. Chromosome.

Images



Western blot analysis of Methyl-Histone H3 (di K4) expression in HeLa cell lysate.

Image not found: 202311/AP90181-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human liver, using Methyl-Histone H3 (di K4) Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.