

TRAF6 Antibody

Rabbit mAb

Catalog # AP90225

Product Information

| | |
|--------------------------|---|
| Application | WB, IHC, IF, ICC, IHF |
| Primary Accession | Q9Y4K3 |
| Reactivity | Rat, Human, Mouse |
| Clonality | Monoclonal |
| Other Names | TNF receptor-associated factor 6; E3 ubiquitin-protein ligase TRAF6; Interleukin-1 signal transducer; RING finger protein 85; TRAF6; RNF85; TRAF 6; TRAF-6; |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 59573 |

Additional Information

| | |
|-------------------------------------|--|
| Dilution | WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human TRAF6 |
| Description | TRAFs (TNF receptor-associated factors) are a family of multifunctional adaptor proteins that bind to surface receptors and recruit additional proteins to form multiprotein signaling complexes capable of promoting cellular responses. Members of the TRAF family share a common carboxy-terminal TRAF domain which mediates interactions with associated proteins; many also contain amino-terminal Zinc/RING finger motifs. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Protein Information

| | |
|-----------------|--|
| Name | TRAF6 |
| Synonyms | RNF85 |
| Function | E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed: 11057907 , PubMed: 18347055 , PubMed: 19465916 , PubMed: 19713527 , PubMed: 27746020 , PubMed: 31620128). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (PubMed: 19675569). Leads to the activation of NF-kappa-B and JUN (PubMed: 16378096 , PubMed: 17135271 , PubMed: 17703191). Seems to also play a role in dendritic cells (DCs) maturation and/or activation (By similarity). |

Represses c-Myb-mediated transactivation, in B-lymphocytes (PubMed:[18093978](#), PubMed:[18758450](#)). Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor (PubMed:[12140561](#), PubMed:[19825828](#), PubMed:[8837778](#)). Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L stimulation (By similarity). Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production (By similarity). Acts as a regulator of the JNK and NF-kappa-B signaling pathways by initiating assembly of heterotypic 'Lys-63'-'Lys-48'-linked branched ubiquitin chains that are then recognized by TAB2: TRAF6 catalyzes initial 'Lys-63'-linked-polyubiquitin chains that are then branched via 'Lys-48'-linked polyubiquitin by HUWE1 (PubMed:[27746020](#)). 'Lys-63'-'Lys-48'-linked branched ubiquitin chains protect 'Lys-63'- linkages from CYLD deubiquitination (PubMed:[27746020](#)). Participates also in the TCR signaling by ubiquitinating LAT (PubMed:[23514740](#), PubMed:[25907557](#)).


Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Nucleus. Lipid droplet {ECO:0000250|UniProtKB:P70196}. Note=Found in the nuclei of some aggressive B-cell lymphoma cell lines as well as in the nuclei of both resting and activated T- and B-lymphocytes. Found in punctate nuclear body protein complexes. Ubiquitination may occur in the cytoplasm and sumoylation in the nucleus. RSAD2/viperin recruits it to the lipid droplet (By similarity).

Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

Images



Western blot analysis of TRAF6 expression in Jurkat cell lysate.

Image not found : 202311/AP90225-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human lung carcinoma, using TRAF6 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.