

Androgen Receptor Antibody

Rabbit mAb

Catalog # AP90368

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	P10275
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	ANDR; Androgen receptor; DHTR; Dihydrotestosterone receptor; NR3C4; SBMA; SMAX1 HYSPI; AR; AIS; TFM; KD;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	99188

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Androgen Receptor
Description	Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Transcription factor activity is modulated by bound coactivator and corepressor proteins. Transcription activation is down-regulated by NROB2. Activated, but not phosphorylated, by HIPK3.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	AR
Synonyms	DHTR, NR3C4
Function	Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues (PubMed: 19022849). Transcription factor activity is modulated by bound coactivator and corepressor proteins like ZBTB7A that recruits NCOR1 and NCOR2 to the androgen response elements/ARE on target genes, negatively regulating androgen receptor signaling and androgen-induced cell proliferation (PubMed: 20812024). Transcription activation is also down-regulated by NROB2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.
Cellular Location	Nucleus. Cytoplasm Note=Detected at the promoter of target genes

(PubMed:25091737) Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.

Tissue Location

[Isoform 2]: Mainly expressed in heart and skeletal muscle.

Images



Western blot analysis of Androgen Receptor expression in Lncap cell lysate.

Image not found : 202311/AP90368-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human prostate, using Androgen Receptor Antibody.

Image not found : 202311/AP90368-IF.jpg

Immunofluorescent analysis of MCF7 cells, using Androgen Receptor Antibody .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.