

CD9 Antibody

Rabbit mAb

Catalog # AP90376

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	P21926
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CD9 antigen; p24; Leukocyte antigen MIC3; Motility-related protein; .MRP-1; Tetraspanin-29; Tspan-29; MIC3; TSPAN29;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	25416

Additional Information

Dilution	WB 1:1000~1:5000 IHC 1:100~1:500 ICC/IF 1:50~1:200 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD9
Description	Involved in platelet activation and aggregation. Regulates paranodal junction formation. Involved in cell adhesion, cell motility and tumor metastasis. Required for sperm-egg fusion.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	CD9 {ECO:0000303 PubMed:1840589, ECO:0000312 HGNC:HGNC:1709}
Function	Integral membrane protein associated with integrins, which regulates different processes, such as sperm-egg fusion, platelet activation and aggregation, and cell adhesion (PubMed: 14575715 , PubMed: 18541721 , PubMed: 8478605). Present at the cell surface of oocytes and plays a key role in sperm-egg fusion, possibly by organizing multiprotein complexes and the morphology of the membrane required for the fusion (By similarity). In myoblasts, associates with CD81 and PTGFRN and inhibits myotube fusion during muscle regeneration (By similarity). In macrophages, associates with CD81 and beta-1 and beta-2 integrins, and prevents macrophage fusion into multinucleated giant cells specialized in ingesting complement-opsonized large particles (PubMed: 12796480). Also prevents the fusion between mononuclear cell progenitors into osteoclasts in charge of bone resorption (By similarity). Acts as a receptor for PSG17 (By similarity). Involved in platelet activation and aggregation (PubMed: 18541721). Regulates paranodal junction formation (By similarity). Involved in cell adhesion, cell motility and tumor

metastasis (PubMed:[7511626](#), PubMed:[8478605](#)).


Cellular Location

Cell membrane; Multi-pass membrane protein. Membrane; Multi-pass membrane protein. Secreted, extracellular exosome {ECO:0000250|UniProtKB:P40240}. Note=Present at the cell surface of oocytes. Accumulates in the adhesion area between the sperm and egg following interaction between IZUMO1 and its receptor IZUMO1R/JUNO {ECO:0000250|UniProtKB:P40240}

Tissue Location

Detected in platelets (at protein level) (PubMed:19640571). Expressed by a variety of hematopoietic and epithelial cells (PubMed:19640571).

Images



Western blot analysis of CD9 expression in (1)HACAT cell lysate ;(2)HepG2 cell lysate.

Image not found : 202311/AP90376-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human tonsil, using CD9 Antibody.

Image not found : 202311/AP90376-IF.jpg

Immunofluorescent analysis of Hela cells, using CD9 Antibody .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.