

CD4 Antibody

Rabbit mAb Catalog # AP90427

Product Information

Application	WB, IHC, IF, FC, ICC, IHF	
Primary Accession	<u>P01730</u>	
Reactivity	Human	
Clonality	Monoclonal	
Other Names	CD4; Leu3; L3T4; Ly4; p55; W3/25;	
Isotype	Rabbit IgG	
Host	Rabbit	
Calculated MW	51111	

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD4
Description	Cluster of Differentiation 4 (CD4) is a glycoprotein composed of an
	amino-terminal extracellular domain (four domains: D1-D4 with Ig-like
	structures), a transmembrane part and a short cytoplasmic tail. CD4 is
	expressed on the surface of T helper cells, regulatory T cells, monocytes,
	macrophages and dendritic cells, and plays an important role in the
	development and activation of T cells. On T cells, CD4 is the co-receptor for
	the T cell receptor (TCR), and these two distinct structures recognize the
	Antigen–Major Histocompatibility Complex (MHC).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium
	azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.

Protein Information

Name

Function

CD4

Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells

	such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T- helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages.
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=Localizes to lipid rafts (PubMed:12517957, PubMed:9168119). Removed from plasma membrane by HIV- 1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope polyprotein gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum
Tissue Location	Highly expressed in T-helper cells. The presence of CD4 is a hallmark of T-helper cells which are specialized in the activation and growth of cytotoxic T-cells, regulation of B cells, or activation of phagocytes. CD4 is also present in other immune cells such as macrophages, dendritic cells or NK cells

Images



Western blot analysis of CD4 expression in THP-1 cell lysate.

Image not found : 202311/AP90427-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human colon, using CD4 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.