

IRF1 Antibody

Rabbit mAb

Catalog # AP90510

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	P10914
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Interferon regulatory factor 1; IRF1; MAR;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	36502

Additional Information

Dilution	WB 1:500~1:1000 IHC 1:50~1:100 ICC/IF 1:50~1:100 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human IRF1
Description	Specifically binds to the upstream regulatory region of type I IFN and IFN-inducible MHC class I genes (the interferon consensus sequence (ICS)) and activates those genes. Acts as a tumor suppressor. Defects in IRF1 are a cause of gastric cancer (GASC) [MIM:613659]; also called gastric cancer intestinal or stomach cancer. Gastric cancer is a malignant disease which starts in the stomach, can spread to the esophagus or the small intestine, and can extend through the stomach wall to nearby lymph nodes and organs. It also can metastasize to other parts of the body.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	IRF1
Function	Transcriptional regulator which displays a remarkable functional diversity in the regulation of cellular responses (PubMed: 15226432 , PubMed: 15509808 , PubMed: 17516545 , PubMed: 17942705 , PubMed: 18497060 , PubMed: 19404407 , PubMed: 19851330 , PubMed: 22367195 , PubMed: 32385160). Regulates transcription of IFN and IFN-inducible genes, host response to viral and bacterial infections, regulation of many genes expressed during hematopoiesis, inflammation, immune responses and cell proliferation and differentiation, regulation of the cell cycle and induction of growth arrest and programmed cell death following DNA damage (PubMed: 15226432 , PubMed: 15509808 , PubMed: 17516545 , PubMed: 17942705 , PubMed: 18497060 , PubMed: 19404407 ,

PubMed:[19851330](#), PubMed:[22367195](#)). Stimulates both innate and acquired immune responses through the activation of specific target genes and can act as a transcriptional activator and repressor regulating target genes by binding to an interferon-stimulated response element (ISRE) in their promoters (PubMed:[15226432](#), PubMed:[15509808](#), PubMed:[17516545](#), PubMed:[17942705](#), PubMed:[18497060](#), PubMed:[19404407](#), PubMed:[19851330](#), PubMed:[21389130](#), PubMed:[22367195](#)). Has an essential role in IFNG- dependent immunity to mycobacteria (PubMed:[36736301](#)). Competes with the transcriptional repressor ZBED2 for binding to a common consensus sequence in gene promoters (PubMed:[32385160](#)). Its target genes for transcriptional activation activity include: genes involved in anti- viral response, such as IFN-alpha/beta, RIGI, TNFSF10/TRAIL, ZBP1, OAS1/2, PIAS1/GBP, EIF2AK2/PKR and RSAD2/viperin; antibacterial response, such as GBP2, GBP5 and NOS2/INOS; anti-proliferative response, such as p53/TP53, LOX and CDKN1A; apoptosis, such as BBC3/PUMA, CASP1, CASP7 and CASP8; immune response, such as IL7, IL12A/B and IL15, PTGS2/COX2 and CYBB; DNA damage responses and DNA repair, such as POLQ/POLH; MHC class I expression, such as TAP1, PSMB9/LMP2, PSME1/PA28A, PSME2/PA28B and B2M and MHC class II expression, such as CIITA; metabolic enzymes, such as ACOD1/IRG1 (PubMed:[15226432](#), PubMed:[15509808](#), PubMed:[17516545](#), PubMed:[17942705](#), PubMed:[18497060](#), PubMed:[19404407](#), PubMed:[19851330](#), PubMed:[22367195](#)). Represses genes involved in anti-proliferative response, such as BIRC5/survivin, CCNB1, CCNE1, CDK1, CDK2 and CDK4 and in immune response, such as FOXP3, IL4, ANXA2 and TLR4 (PubMed:[18641303](#), PubMed:[22200613](#)). Stimulates p53/TP53-dependent transcription through enhanced recruitment of EP300 leading to increased acetylation of p53/TP53 (PubMed:[15509808](#), PubMed:[18084608](#)). Plays an important role in immune response directly affecting NK maturation and activity, macrophage production of IL12, Th1 development and maturation of CD8+ T-cells (PubMed:[11244049](#), PubMed:[11846971](#), PubMed:[11846974](#), PubMed:[16932750](#)). Also implicated in the differentiation and maturation of dendritic cells and in the suppression of regulatory T (Treg) cells development (PubMed:[11244049](#), PubMed:[11846971](#), PubMed:[11846974](#), PubMed:[16932750](#)). Acts as a tumor suppressor and plays a role not only in antagonism of tumor cell growth but also in stimulating an immune response against tumor cells (PubMed:[20049431](#)).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:P15314}.
Note=MYD88-associated IRF1 migrates into the nucleus more efficiently than non-MYD88-associated IRF1 {ECO:0000250|UniProtKB:P15314}

Images



Western blot analysis of IRF1 expression in Jurkat cell lysate.

Image not found : 202311/AP90510-IHC.jpg

Immunohistochemical analysis of paraffin-embedded rat kidney, using IRF1 Antibody.

Image not found : 202311/AP90510-IF.jpg

Immunofluorescent analysis of Hela cells, using IRF1 Antibody .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.