

Phospho-GATA3 (S308) Antibody

Rabbit mAb Catalog # AP90511

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IP <u>P23771</u> Human Monoclonal GATA-binding factor 3; Trans-acting T-cell specific transcription factor GATA-3; GATA3;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	47916

Additional Information

Dilution Purification Immunogen	WB 1:500~1:2000 IHC 1:50~1:100 IP 1:50 Affinity-chromatography A synthesized peptide derived from human Phospho-GATA3 (S308)
Description	Transcriptional activator which binds to the enhancer of the T-cell receptor
Description	alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	GATA3
Function	Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses. Positively regulates ASB2 expression (By similarity). Coordinates macrophage transcriptional activation and UCP2-dependent metabolic reprogramming in response to IL33. Upon tissue injury, acts downstream of IL33 signaling to drive differentiation of inflammation-resolving alternatively activated macrophages.
Cellular Location	Nucleus.
Tissue Location	T-cells and endothelial cells.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.