

ASH2L Antibody

Rabbit mAb

Catalog # AP90529

Product Information

| | |
|--------------------------|---|
| Application | WB, IHC, IF, ICC, IHF |
| Primary Accession | Q9UBL3 |
| Reactivity | Rat, Human, Mouse |
| Clonality | Monoclonal |
| Other Names | ASH 2; ASH2; ASH2 LIKE; ASH2 like protein; ASH2-like protein; Ash2L; ASH2L1; ASH2L2; Bre 2; Bre2; |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 68723 |

Additional Information

| | |
|-------------------------------------|--|
| Dilution | WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human ASH2L |
| Description | Component of the Set1/Ash2 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3, but not if the neighboring 'Lys-9' residue is already methylated. As part of the MLL1/MLL complex it is involved in methylation and dimethylation at 'Lys-4' of histone H3. May function as a transcriptional regulator. May play a role in hematopoiesis. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Protein Information

| | |
|-----------------|---|
| Name | ASH2L |
| Synonyms | ASH2L1 |
| Function | Transcriptional regulator (PubMed: 12670868). Component or associated component of some histone methyltransferase complexes which regulates transcription through recruitment of those complexes to gene promoters (PubMed: 19131338). Component of the Set1/Ash2 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3, but not if the neighboring 'Lys-9' residue is already methylated (PubMed: 19556245). As part of the MLL1/MLL complex it is involved in methylation and dimethylation at 'Lys-4' of histone H3 (PubMed: 19556245). May play a role in hematopoiesis (PubMed: 12670868). In association with RBBP5 and WDR5, stimulates the histone methyltransferase activities of |

KMT2A, KMT2B, KMT2C, KMT2D, SETD1A and SETD1B (PubMed:[21220120](#), PubMed:[22266653](#)).


Cellular Location

Nucleus.

Tissue Location

Ubiquitously expressed. Predominantly expressed in adult heart and testis and fetal lung and liver, with barely detectable expression in adult lung, liver, kidney, prostate, and peripheral leukocytes.

Images



Western blot analysis of ASH2L expression in K562 cell lysate.

Image not found : 202311/AP90529-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human kidney, using ASH2L Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.