10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Histone H3.3 Antibody

Rabbit mAb
Catalog # AP90553

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	P84243
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	H3.3; H3.3A; H33_HUMAN ; H3F3; Histone H3.3 ; H3 histone family 3A; H3 histone family 3B
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	15328

Additional Information

Dilution	WB 1:500~1:1000 IHC 1:500~1:1000 ICC/IF 1:500~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Histone H3.3
Description	Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Protein Information

Name	H3-3A (HGNC:4764)
Synonyms	H3.3A, H3F3, H3F3A
Function	Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription


regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location Nucleus. Chromosome

Images



Western blot analysis of Histone H3.3 expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate.



Immunohistochemical analysis of paraffin-embedded mouse stomach, using Histone H3.3 Antibody .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.