

Cyclin E2 Antibody

Rabbit mAb Catalog # AP90570

Product Information

Application	WB, IHC, IF, ICC, IP, IHF
Primary Accession	<u>O96020</u>
Reactivity	Human
Clonality	Monoclonal
Other Names	CGE2; CYCE2; cyclin E2; G1/S-Specific cyclin E2;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	46757

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human Cyclin E2 The human Cyclin E2 gene encodes a 404 amino acid protein that is most closely related to Cyclin E. Cyclin E2 mRNA levels peaks at the G1 / S transition. Cyclin E2 associates with Cdk2 in a functional kinase complex that is inhibited by both p27 (Kip1) and p21 (Cip1). Cyclin E2 / Cdk2 phosphorylates histone H1 in vitro. G1 cyclin E controls the initiation of DNA synthesis by activating CDK2. Abnormally high levels of cyclin E expression have frequently
Storage Condition and Buffer	been observed in human cancers. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	CCNE2
Function	Essential for the control of the cell cycle at the late G1 and early S phase.
Cellular Location	Nucleus.
Tissue Location	According to PubMed:9858585, highest levels of expression in adult testis, thymus and brain. Lower levels in placenta, spleen and colon. Consistently elevated levels in tumor-derived cells compared to non-transformed proliferating cells. According to PubMed:9840927: low levels in thymus, prostate, brain, skeletal muscle, and kidney. Elevated levels in lung. According to PubMed:9840943 highly expressed in testis, placenta, thymus and brain. In a lesser extent in small intestine and colon.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.