

Prion Protein Antibody

Rabbit mAb

Catalog # AP90618

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	P04156
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CJD; GSS; PrP; PRNP; ASCR; KURU; PRIP; PrPc; CD230; AltPrP; Prion protein; PrP27-30; PrP33-35C;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	27661

Additional Information


Dilution	WB 1:1000~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Prion Protein
Description	The PRNP gene encodes the major prion protein (PrP, CD230), a widely-expressed glycoprotein expressed at high levels in the central nervous system. While the typical cellular function of PrP is not well defined, it is a putative antioxidant and a metal-binding protein that may be involved in signal transduction.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	PRNP
Synonyms	ALTPRP, PRIP, PRP
Function	Its primary physiological function is unclear. May play a role in neuronal development and synaptic plasticity. May be required for neuronal myelin sheath maintenance. May promote myelin homeostasis through acting as an agonist for ADGRG6 receptor. May play a role in iron uptake and iron homeostasis. Soluble oligomers are toxic to cultured neuroblastoma cells and induce apoptosis (in vitro) (By similarity). Association with GPC1 (via its heparan sulfate chains) targets PRNP to lipid rafts. Also provides Cu(2+) or Zn(2+) for the ascorbate-mediated GPC1 deaminase degradation of its heparan sulfate side chains (By similarity).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor. Golgi apparatus

{ECO:0000250|UniProtKB:P04925}. Note=Targeted to lipid rafts via association with the heparan sulfate chains of GPC1. Colocalizes, in the presence of Cu(2+), to vesicles in para- and perinuclear regions, where both proteins undergo internalization. Heparin displaces PRNP from lipid rafts and promotes endocytosis.

Images



Western blot analysis of Prion Protein expression in Hunam fetal brain lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.