

PI3 Kinase p110 beta Antibody

Rabbit mAb Catalog # AP90623

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, FC, IP <u>P42338</u> Human Monoclonal PIK3CB; DKFZp779K1237; MGC133043; PI3K; PI3KCB; PI3Kbeta; PIK3C1; p110-BETA;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	122762

Additional Information

Dilution	WB 1:500~1:2000 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PI3 Kinase p110 beta
Description	Phosphoinositide 3-kinase (PI3K) catalyzes the production of
	phosphatidylinositol-3,4,5-triphosphate by phosphorylating
	phosphatidylinositol (PI), phosphatidylinositol-4-phosphate (PIP) and
	phosphatidylinositol-4,5-bisphosphate (PIP2). Growth factors and hormones
	trigger this phosphorylation event, which in turn coordinates cell growth, cell
	cycle entry, cell migration, and cell survival.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium
	azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.

Protein Information

Name	PIK3CB
Synonyms	PIK3C1
Function	Phosphoinositide-3-kinase (PI3K) phosphorylates phosphatidylinositol derivatives at position 3 of the inositol ring to produce 3-phosphoinositides (PubMed: <u>15135396</u>). Uses ATP and PtdIns(4,5)P2 (phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3) (PubMed: <u>15135396</u>). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDPK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Involved in the activation of AKT1 upon stimulation by G- protein coupled receptors (GPCRs) ligands such as CXCL12, sphingosine 1-phosphate, and lysophosphatidic acid. May also act downstream receptor tyrosine

	kinases. Required in different signaling pathways for stable platelet adhesion and aggregation. Plays a role in platelet activation signaling triggered by GPCRs, alpha-IIb/beta-3 integrins (ITGA2B/ ITGB3) and ITAM (immunoreceptor tyrosine-based activation motif)-bearing receptors such as GP6. Regulates the strength of adhesion of ITGA2B/ ITGB3 activated receptors necessary for the cellular transmission of contractile forces. Required for platelet aggregation induced by F2 (thrombin) and thromboxane A2 (TXA2). Has a role in cell survival. May have a role in cell migration. Involved in the early stage of autophagosome formation. Modulates the intracellular level of PtdIns3P (phosphatidylinositol 3-phosphate) and activates PIK3C3 kinase activity. May act as a scaffold, independently of its lipid kinase activity to positively regulate autophagy. May have a role in insulin signaling as scaffolding protein in which the lipid kinase activity is not required. May have a kinase-independent function in regulating cell proliferation and in clathrin-mediated endocytosis. Mediator of oncogenic signal in cell lines lacking PTEN. The lipid kinase activity is necessary for its role in oncogenic transformation. Required for the growth of ERBB2 and RAS driven tumors. Also has a protein kinase activity showing autophosphorylation (PubMed:12502714).
Cellular Location	Cytoplasm. Nucleus. Note=Interaction with PIK3R2 is required for nuclear localization and export
Tissue Location	Expressed ubiquitously.

Images



Western blot analysis of PI3 Kinase p110 beta expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.