10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-c-Myc (T58) Antibody

Rabbit mAb
Catalog # AP90633

Product Information

Application	WB, IF, FC, ICC
Primary Accession	P01106
Reactivity	Human
Clonality	Monoclonal
Other Names	MRTL; MYC; Myc proto-oncogene protein; c-myc;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	50565

Additional Information


Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Phospho-c-Myc (T58)
Description	Myc a proto-oncogenic transcription factor that plays a role in cell proliferation, apoptosis and in the development of human tumors. Seems to activate the transcription of growth-related genes.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	MYC
Synonyms	BHLHE39
Function	Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed:24940000, PubMed:25956029). Activates the transcription of growth-related genes (PubMed:24940000, PubMed:25956029). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:24940000, PubMed:25956029). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:20010808).

Cellular Location	Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Chromosome. Note=Association with chromatin is reduced by hyperphosphorylation (PubMed:30158517) Localization to the nucleolus is dependent on HEATR1 (PubMed:38225354)

Images



Western blot analysis of Phospho-c-Myc (T58) expression in (1) HeLa cell lysate; (2) HeLa cell lysate treated with Calyculin A and Okadaic Acid.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.