

VAV2 Antibody

Rabbit mAb Catalog # AP90715

Product Information

| Application | WB, IHC, IF, ICC, IP, IHF |
|-------------------|------------------------------|
| Primary Accession | <u>P52735</u> |
| Reactivity | Rat, Human, Mouse |
| Clonality | Monoclonal |
| Other Names | VAV2; Vav 2 oncogene; VAV-2; |
| lsotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 101289 |

Additional Information

| Dilution Purification Immunogen | WB 1:5000~1:20000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:100 Affinity-chromatography A synthesized peptide derived from human VAV2 |
|---------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Description | Vav proteins belong to the Dbl family of guanine nucleotide exchange factors (GEFs) for Rho/Rac small GTPases. The three identified mammalian Vav proteins (Vav1, Vav2 and Vav3) differ in their expression. Vav1 is expressed only in hematopoietic cells and is involved in the formation of the immune synapse. Vav2 and Vav3 are more ubiguitously expressed. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Protein Information

| Name | VAV2 |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Function | Guanine nucleotide exchange factor for the Rho family of Ras- related GTPases. Plays an important role in angiogenesis. Its recruitment by phosphorylated EPHA2 is critical for EFNA1-induced RAC1 GTPase activation and vascular endothelial cell migration and assembly (By similarity). |
| Tissue Location | Widely expressed. |
| Images | |

Western blot analysis of VAV2 expression in 293T cell lysate.



Image not found : 202311/AP90715-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human breast cancer, using VAV2 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.