10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Caspase-8 Antibody

Rabbit mAb
Catalog # AP90751

## Product Information

| | |
|---|---|
| **Application** | WB, IHC, IF, ICC, IHF |
| **Primary Accession** | Q14790 |
| **Reactivity** | Rat, Human, Mouse |
| **Clonality** | Monoclonal |
| **Other Names** | Caspase 8; CASP-8; Apoptotic cysteine protease; Apoptotic protease Mch-5; FADD-homologous ICE/ced-3-like protease; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; MACH; Caspase-8 subunit p18; CAP4; |
| **Isotype** | Rabbit IgG |
| **Host** | Rabbit |
| **Calculated MW** | 55391 |

## Additional Information

| | |
|---|---|
| **Dilution** | WB 1:1000~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 |
| **Purification** | Affinity-chromatography |
| **Immunogen** | A synthesized peptide derived from human Caspase-8 |
| **Description** | Caspases are a family of cytosolic aspartate specific cysteine proteases. Involved in the activation cascade of caspases responsible for apoptosis execution. Activated caspase-8 cleaves and activates downstream effector caspases such as caspase-1, -3, -6, and -7. |
| **Storage Condition and Buffer** | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

## Protein Information

| | |
|---|---|
| **Name** | CASP8 {ECO:0000303\|PubMed:9931493, ECO:0000312\|HGNC:HGNC:1509} |
| **Function** | Thiol protease that plays a key role in programmed cell death by acting as a molecular switch for apoptosis, necroptosis and pyroptosis, and is required to prevent tissue damage during embryonic development and adulthood (PubMed:23516580, PubMed:35338844, PubMed:35446120, PubMed:8681376, PubMed:8681377, PubMed:8962078, PubMed:9006941, PubMed:9184224). Initiator protease that induces extrinsic apoptosis by mediating cleavage and activation of effector caspases responsible for FAS/CD95-mediated and TNFRSF1A-induced cell death (PubMed:23516580, PubMed:35338844, PubMed:35446120, PubMed:8681376, PubMed:8681377, PubMed:8962078, PubMed:9006941, PubMed:9184224). Cleaves and activates effector caspases CASP3, CASP4, CASP6, CASP7, CASP9 and CASP10 (PubMed:16916640, PubMed:8962078, PubMed:9006941). Binding to the adapter molecule FADD recruits it to either receptor FAS/TNFRSF6 or |

TNFRSF1A (PubMed:8681376, PubMed:8681377). The resulting aggregate called the death-inducing signaling complex (DISC) performs CASP8 proteolytic activation (PubMed:9184224). The active dimeric enzyme is then liberated from the DISC and free to activate downstream apoptotic proteases (PubMed:9184224). Proteolytic fragments of the N-terminal propeptide (termed CAP3, CAP5 and CAP6) are likely retained in the DISC (PubMed:9184224). Also cleaves and activates BID, thereby promoting cytochrome C release from mitochrondria (By similarity). In addition to extrinsic apoptosis, also acts as a negative regulator of necroptosis: acts by cleaving RIPK1 at 'Asp-324', which is crucial to inhibit RIPK1 kinase activity, limiting TNF-induced apoptosis, necroptosis and inflammatory response (PubMed:31827280, PubMed:31827281). Also able to initiate pyroptosis by mediating cleavage and activation of gasdermin-C and -D (GSDMC and GSDMD, respectively): gasdermin cleavage promotes release of the N-terminal moiety that binds to membranes and forms pores, triggering pyroptosis (PubMed:32929201, PubMed:34012073). Initiates pyroptosis following inactivation of MAP3K7/TAK1 (By similarity). Also acts as a regulator of innate immunity by mediating cleavage and inactivation of N4BP1 downstream of TLR3 or TLR4, thereby promoting cytokine production (By similarity). May participate in the Granzyme B (GZMB) cell death pathways (PubMed:8755496). Cleaves PARP1 and PARP2 (PubMed:8681376). Independent of its protease activity, promotes cell migration following phosphorylation at Tyr-380 (PubMed:18216014, PubMed:27109099).


**Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q9JHX4}. Nucleus {ECO:0000250|UniProtKB:Q9JHX4}. Cell projection, lamellipodium. Note=Recruitment to lamellipodia of migrating cells is enhanced by phosphorylation at Tyr-380

**Tissue Location**

Isoform 1, isoform 5 and isoform 7 are expressed in a wide variety of tissues. Highest expression in peripheral blood leukocytes, spleen, thymus and liver. Barely detectable in brain, testis and skeletal muscle

## Images




Western blot analysis of Caspase-8 expression in HeLa cell lysate.

Image not found : 202311/AP90751-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human uterus, using Caspase-8 Antibody.

Image not found : 202311/AP90751-IF.jpg

Immunofluorescent analysis of Hela cells, using Caspase-8 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.