

# Histone H3 (di methyl K9) Antibody

Rabbit mAb Catalog # AP90762

## **Product Information**

Application	WB, IF, ICC, IP
Primary Accession	<u>P68431</u>
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Histone H3.1; Histone H3; HIST1H3A; H3K9me2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	15404


## **Additional Information**

Dilution Purification Immunogen	WB 1:500~1:1000 ICC/IF 1:50~1:200 ChIP Affinity-chromatography A synthesized peptide derived from human Histone H3
Description	Belongs to the histone H3 family. Play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### **Protein Information**

Name	H3C1 ( <u>HGNC:4766</u> )
Synonyms	H3FA, HIST1H3A
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.

### Images



Western blot analysis of Histone H3 (di methyl K9) expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.