

Caspase-3 p12 Antibody

Rabbit mAb Catalog # AP90784

Product Information

Application WB, IHC, IF, ICC, IP, IHF

Primary Accession P42574

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names Apopain; CASP-3; CASP3; Caspase-3 subunit p12; CPP-32; Cysteine protease

CPP32; Protein Yama; SCA-1; SREBP cleavage activity 1;

IsotypeRabbit IgGHostRabbitCalculated MW31608

Additional Information

Dilution WB 1:500~1:1000 IHC 1:100~1:500 ICC/IF 1:50~1:200

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human Caspase-3 p12

Description Caspase-3 is a critical executioner of apoptosis, as it is either partially or

totally responsible for the proteolytic cleavage of many key proteins, such as the nuclear enzyme poly (ADP-ribose) polymerase (PARP). Involved in the activation cascade of caspases responsible for apoptosis execution.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name CASP3

Synonyms CPP32 {ECO:0000303 | PubMed:7983002}

Function Thiol protease that acts as a major effector caspase involved in the

execution phase of apoptosis (PubMed: 18723680, PubMed: 20566630,

PubMed: 23650375, PubMed: 35338844, PubMed: 35446120,

PubMed: <u>7596430</u>). Following cleavage and activation by initiator caspases (CASP8, CASP9 and/or CASP10), mediates execution of apoptosis by catalyzing

cleavage of many proteins (PubMed: 18723680, PubMed: 20566630, PubMed: 23650375, PubMed: 7596430). At the onset of apoptosis, it proteolytically cleaves poly(ADP-ribose) polymerase PARP1 at a '216-Asp-|-Gly-217' bond (PubMed: 10497198, PubMed: 16374543,

PubMed:<u>7596430</u>, PubMed:<u>7774019</u>). Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain (By similarity). Cleaves

and activates caspase-6, -7 and -9 (CASP6, CASP7 and CASP9, respectively) (PubMed: <u>7596430</u>). Cleaves and inactivates interleukin-18 (IL18) (PubMed:<u>37993714</u>, PubMed:<u>9334240</u>). Involved in the cleavage of huntingtin (PubMed:8696339). Triggers cell adhesion in sympathetic neurons through RET cleavage (PubMed:21357690). Cleaves and inhibits serine/threonine-protein kinase AKT1 in response to oxidative stress (PubMed:23152800). Acts as an inhibitor of type I interferon production during virus-induced apoptosis by mediating cleavage of antiviral proteins CGAS, IRF3 and MAVS, thereby preventing cytokine overproduction (PubMed:30878284). Also involved in pyroptosis by mediating cleavage and activation of gasdermin-E (GSDME) (PubMed:35338844, PubMed:35446120). Cleaves XRCC4 and phospholipid scramblase proteins XKR4, XKR8 and XKR9, leading to promote phosphatidylserine exposure on apoptotic cell surface (PubMed:23845944, PubMed:33725486). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed:36758104, PubMed:36758106).


Cellular Location

Cytoplasm.

Tissue Location

Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.

Images



Western blot analysis of Caspase-3 p12 expression in HeLa cell treated with staurosporine lysate.

Image not found: 202311/AP90784-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human kidney, using Caspase-3 p12 Antibody.

Image not found: 202311/AP90784-IF.jpg

Immunofluorescent analysis of HeLa cells, using Caspase-3 p12 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.