

PDPK1 Antibody

Rabbit mAb

Catalog # AP90834

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	O15530
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	3-phosphoinositide dependent protein kinase-1; EC 2.7.11.1; hPDK1; kinase PDK1; mPDK1; PDPK1; PkB kinase; Protein kinase B kinase;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	63152

Additional Information

Dilution	WB 1:1000~1:2000 IHC 1:100~1:500 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PDPK1
Description	Phosphoinositide-dependent protein kinase 1 (PDK1) plays a central role in many signal transduction pathways including the activation of Akt and the PKC isoenzymes p70 S6 kinase and RSK. Through its effects on these kinases, PDK1 is involved in the regulation of a wide variety of processes, including cell proliferation, differentiation and apoptosis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	PDPK1
Synonyms	PDK1
Function	Serine/threonine kinase which acts as a master kinase, phosphorylating and activating a subgroup of the AGC family of protein kinases (PubMed: 10226025 , PubMed: 10480933 , PubMed: 10995762 , PubMed: 12167717 , PubMed: 14585963 , PubMed: 14604990 , PubMed: 16207722 , PubMed: 16251192 , PubMed: 17327236 , PubMed: 17371830 , PubMed: 18835241 , PubMed: 9094314 , PubMed: 9368760 , PubMed: 9445476 , PubMed: 9445477 , PubMed: 9707564 , PubMed: 9768361). Its targets include: protein kinase B (PKB/AKT1, PKB/AKT2, PKB/AKT3), p70 ribosomal protein S6 kinase (RPS6KB1), p90 ribosomal protein S6 kinase (RPS6KA1, RPS6KA2 and RPS6KA3), cyclic AMP-dependent protein kinase (PRKACA), protein kinase C (PRKCD and PRK CZ), serum and

glucocorticoid-inducible kinase (SGK1, SGK2 and SGK3), p21-activated kinase-1 (PAK1), TSSK3, protein kinase PKN (PKN1 and PKN2) (PubMed:[10226025](#), PubMed:[10480933](#), PubMed:[10995762](#), PubMed:[12167717](#), PubMed:[14585963](#), PubMed:[14604990](#), PubMed:[16207722](#), PubMed:[16251192](#), PubMed:[17327236](#), PubMed:[17371830](#), PubMed:[18835241](#), PubMed:[9094314](#), PubMed:[9368760](#), PubMed:[9445476](#), PubMed:[9707564](#), PubMed:[9768361](#)). Plays a central role in the transduction of signals from insulin by providing the activating phosphorylation to PKB/AKT1, thus propagating the signal to downstream targets controlling cell proliferation and survival, as well as glucose and amino acid uptake and storage (PubMed:[10226025](#), PubMed:[12167717](#), PubMed:[9094314](#)). Negatively regulates the TGF-beta-induced signaling by: modulating the association of SMAD3 and SMAD7 with TGF-beta receptor, phosphorylating SMAD2, SMAD3, SMAD4 and SMAD7, preventing the nuclear translocation of SMAD3 and SMAD4 and the translocation of SMAD7 from the nucleus to the cytoplasm in response to TGF-beta (PubMed:[17327236](#)). Activates PPAR γ transcriptional activity and promotes adipocyte differentiation (By similarity). Activates the NF-kappa-B pathway via phosphorylation of IKK β (PubMed:[16207722](#)). The tyrosine phosphorylated form is crucial for the regulation of focal adhesions by angiotensin II (PubMed:[14585963](#)). Controls proliferation, survival, and growth of developing pancreatic cells (By similarity). Participates in the regulation of Ca²⁺ entry and Ca²⁺-activated K⁺ channels of mast cells (By similarity). Essential for the motility of vascular endothelial cells (ECs) and is involved in the regulation of their chemotaxis (PubMed:[17371830](#)). Plays a critical role in cardiac homeostasis by serving as a dual effector for cell survival and beta-adrenergic response (By similarity). Plays an important role during thymocyte development by regulating the expression of key nutrient receptors on the surface of pre-T cells and mediating Notch-induced cell growth and proliferative responses (By similarity). Provides negative feedback inhibition to toll-like receptor-mediated NF-kappa-B activation in macrophages (By similarity).

Cellular Location

Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein. Cell junction, focal adhesion. Note=Tyrosine phosphorylation seems to occur only at the cell membrane. Translocates to the cell membrane following insulin stimulation by a mechanism that involves binding to GRB14 and INSR. SRC and HSP90 promote its localization to the cell membrane. Its nuclear localization is dependent on its association with PTPN6 and its phosphorylation at Ser- 396. Restricted to the nucleus in neuronal cells while in non-neuronal cells it is found in the cytoplasm. The Ser-241 phosphorylated form is distributed along the perinuclear region in neuronal cells while in non-neuronal cells it is found in both the nucleus and the cytoplasm IGF1 transiently increases phosphorylation at Ser-241 of neuronal PDK1, resulting in its translocation to other cellular compartments The tyrosine-phosphorylated form colocalizes with PTK2B in focal adhesions after angiotensin II stimulation

Tissue Location

Appears to be expressed ubiquitously. The Tyr-9 phosphorylated form is markedly increased in diseased tissue compared with normal tissue from lung, liver, colon and breast

Images

Western blot analysis of PDK1 expression in HEK293 cell lysate.




Image not found : 202311/AP90834-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human kidney, using PDPK1 Antibody.

Image not found : 202311/AP90834-IF.jpg

Immunofluorescent analysis of PC-12 cells, using PDPK1 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.