

# TRAF2 Antibody

Rabbit mAb

Catalog # AP90841

## Product Information

<b>Application</b>	WB, IHC, IF, FC, ICC, IP, IHF
<b>Primary Accession</b>	<a href="#">Q12933</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	TNF receptor-associated factor 2; E3 ubiquitin-protein ligase TRAF2; Tumor necrosis factor type 2 receptor-associated protein 3; TRAF2; TRAP3;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	55859

## Additional Information

<b>Dilution</b>	WB 1:500~1:1000 ICC/IF 1:50~1:200 IP 1:40 FC 1:50
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human TRAF2
<b>Description</b>	TRAFs (TNF receptor-associated factors) are a family of multifunctional adaptor proteins that bind to surface receptors and recruit additional proteins to form multiprotein signaling complexes capable of promoting cellular responses. Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR. In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	TRAF2 {ECO:0000303   PubMed:28489822, ECO:0000312   HGNC:HGNC:12032}
<b>Function</b>	E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed: <a href="#">10346818</a> , PubMed: <a href="#">11784851</a> , PubMed: <a href="#">12917689</a> , PubMed: <a href="#">15383523</a> , PubMed: <a href="#">18981220</a> , PubMed: <a href="#">19150425</a> , PubMed: <a href="#">19810754</a> , PubMed: <a href="#">19918265</a> , PubMed: <a href="#">19937093</a> , PubMed: <a href="#">20047764</a> , PubMed: <a href="#">20064526</a> , PubMed: <a href="#">20385093</a> , PubMed: <a href="#">20577214</a> , PubMed: <a href="#">22212761</a> ). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed: <a href="#">23453969</a> , PubMed: <a href="#">28489822</a> ). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed: <a href="#">15383523</a> , PubMed: <a href="#">18981220</a> ). Regulates BIRC2


and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:[11907583](#), PubMed:[19506082](#)). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:[15121867](#)). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE (PubMed:[23453969](#)). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:[28489822](#)). Required for normal antibody isotype switching from IgM to IgG (By similarity).

#### Cellular Location

Cytoplasm

## Images

---



Western blot analysis of TRAF2 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.