

Stromal interaction molecule 1 Antibody

Rabbit mAb

Catalog # AP90887

Product Information

Application	WB, IHC, IP
Primary Accession	Q13586
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	GOK; SIM; STIM 1; Stim1 stromal interaction molecule 1; STIM1L; Stromal interaction molecule 1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	77423

Additional Information

Dilution	WB 1:1000~1:2000 IHC 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Stromal interaction molecule 1
Description	STIM1 is a potential tumor suppressor; defects in STIM1 may cause rhabdomyosarcoma and rhabdoid tumors. STIM1 can either homodimerize or form heterodimers with STIM2. STIM2 possesses a high sequence identity to STIM1 and can function as an inhibitor of STIM1-mediated plasma membrane store-operated Ca^{2+} entry. However, further investigation is required to elucidate the true physiological function of STIM2.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	STIM1
Synonyms	GOK {ECO:0000303 PubMed:9377559}
Function	Acts as a Ca^{2+} sensor that gates two major inward rectifying Ca^{2+} channels at the plasma membrane: Ca^{2+} release- activated Ca^{2+} (CRAC) channels and arachidonate-regulated Ca^{2+} - selective (ARC) channels (PubMed: 15866891 , PubMed: 16005298 , PubMed: 16208375 , PubMed: 16537481 , PubMed: 16733527 , PubMed: 16766533 , PubMed: 16807233 , PubMed: 18854159 , PubMed: 19182790 , PubMed: 19249086 , PubMed: 19622606 , PubMed: 19706554 , PubMed: 22464749 , PubMed: 24069340 , PubMed: 24351972 , PubMed: 24591628 , PubMed: 25326555 , PubMed: 26322679 , PubMed: 28219928 , PubMed: 32415068). Plays a role in mediating store-

operated Ca^{2+} entry (SOCE), a Ca^{2+} influx following depletion of intracellular Ca^{2+} stores. Upon Ca^{2+} depletion, translocates from the endoplasmic reticulum to the plasma membrane where it activates CRAC channel pore-forming subunits ORA1, ORA2 and ORAI3 to generate sustained and oscillatory Ca^{2+} entry (PubMed:[16208375](#), PubMed:[16537481](#), PubMed:[32415068](#)). Involved in enamel formation (PubMed:[24621671](#)).


Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton. Sarcoplasmic reticulum. Note=Translocates from the endoplasmic reticulum to the cell membrane in response to a depletion of intracellular calcium and is detected at punctae corresponding to junctions between the endoplasmic reticulum and the cell membrane (PubMed:16005298, PubMed:16208375, PubMed:18854159, PubMed:19182790, PubMed:19249086). Associated with the microtubule network at the growing distal tip of microtubules (PubMed:19632184). Colocalizes with ORAI1 at the cell membrane (PubMed:27185316). Colocalizes preferentially with CASQ1 at endoplasmic reticulum in response to a depletion of intracellular calcium (PubMed:27185316)

Tissue Location

Ubiquitously expressed in various human primary cells and tumor cell lines.

Images



Western blot analysis of Stromal interaction molecule 1 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.