

HDAC6 Antibody

Rabbit mAb

Catalog # AP90889

Product Information

Application	WB, IHC, IF, ICC, IP, IHF
Primary Accession	Q9UBN7
Reactivity	Human
Clonality	Monoclonal
Other Names	HD 6; HDAC 6; Histone deacetylase 6 (HD6); JM 21;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	131419

Additional Information

Dilution	WB 1:5000~1:20000 IHC 1:50~1:200 ICC/IF 1:100~1:500 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HDAC6
Description	Involved in the regulation of many cellular processes, including cell migration, immune synapse formation, viral infection, and degradation of misfolded proteins. HDAC6 binds to both poly-ubiquitinated misfolded proteins and dynein motors, facilitating the transport of misfolded proteins to the aggresome. Required for subsequent recruitment of the autophagic machinery and clearance of aggresomes from the cell. Plays a key role in the protection against the deleterious effects of pathological protein aggregation that occurs in various diseases.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	HDAC6 {ECO:0000303 PubMed:10220385, ECO:0000312 HGNC:HGNC:14064}
Function	Deacetylates a wide range of non-histone substrates (PubMed: 12024216 , PubMed: 18606987 , PubMed: 20308065 , PubMed: 24882211 , PubMed: 26246421 , PubMed: 30538141 , PubMed: 31857589 , PubMed: 30770470 , PubMed: 38534334 , PubMed: 39567688). Plays a central role in microtubule- dependent cell motility by mediating deacetylation of tubulin (PubMed: 12024216 , PubMed: 20308065 , PubMed: 26246421). Required for cilia disassembly via deacetylation of alpha-tubulin (PubMed: 17604723 , PubMed: 26246421). Alpha-tubulin deacetylation results in destabilization of dynamic microtubules (By similarity). Promotes deacetylation of CTTN, leading to actin polymerization, promotion of autophagosome-lysosome


fusion and completion of autophagy (PubMed:[30538141](#)). Deacetylates SQSTM1 (PubMed:[31857589](#)). Deacetylates peroxiredoxins PRDX1 and PRDX2, decreasing their reducing activity (PubMed:[18606987](#)). Deacetylates antiviral protein RIGI in the presence of viral mRNAs which is required for viral RNA detection by RIGI (By similarity). Sequentially deacetylates and polyubiquitinates DNA mismatch repair protein MSH2 which leads to MSH2 degradation, reducing cellular sensitivity to DNA-damaging agents and decreasing cellular DNA mismatch repair activities (PubMed:[24882211](#)). Deacetylates DNA mismatch repair protein MLH1 which prevents recruitment of the MutL alpha complex (formed by the MLH1-PMS2 heterodimer) to the MutS alpha complex (formed by the MSH2-MSH6 heterodimer), leading to tolerance of DNA damage (PubMed:[30770470](#)). Deacetylates RHOT1/MIRO1 which blocks mitochondrial transport and mediates axon growth inhibition (By similarity). Deacetylates transcription factor SP1 which leads to increased expression of ENG, positively regulating angiogenesis (PubMed:[38534334](#)). Deacetylates KHDRBS1/SAM68 which regulates alternative splicing by inhibiting the inclusion of CD44 alternate exons (PubMed:[26080397](#)). Acts as a valine sensor by binding to valine through the primate-specific SE14 repeat region (PubMed:[39567688](#)). In valine deprivation conditions, translocates from the cytoplasm to the nucleus where it deacetylates TET2 which promotes TET2-dependent DNA demethylation, leading to DNA damage (PubMed:[39567688](#)). Promotes odontoblast differentiation following IPO7-mediated nuclear import and subsequent repression of RUNX2 expression (By similarity). In addition to its protein deacetylase activity, plays a key role in the degradation of misfolded proteins: when misfolded proteins are too abundant to be degraded by the chaperone refolding system and the ubiquitin-proteasome, mediates the transport of misfolded proteins to a cytoplasmic juxtanuclear structure called aggresome (PubMed:[17846173](#)). Probably acts as an adapter that recognizes polyubiquitinated misfolded proteins and targets them to the aggresome, facilitating their clearance by autophagy (PubMed:[17846173](#)). Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer (PubMed:[24413532](#)).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Nucleus. Perikaryon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, axon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body Note=Mainly cytoplasmic where it is associated with microtubules (PubMed:12024216). Can shuttle between the cytoplasm and the nucleus (PubMed:39567688). Retained in the cytoplasm by binding to valine via the primate-specific SE14 repeat region while valine deprivation induces nuclear localization (PubMed:39567688). Found exclusively in the cytoplasm in proliferative cells with a fraction found in the nucleus during differentiation (By similarity). May translocate to the nucleus following DNA damage (PubMed:30770470) {ECO:0000250|UniProtKB:Q9Z2V5, ECO:0000269|PubMed:12024216, ECO:0000269|PubMed:30770470, ECO:0000269|PubMed:39567688}

Images

Western blot analysis of HDAC6 expression in HeLa cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.