

## Cdc23/APC8 Antibody

Rabbit mAb Catalog # AP90946

## **Product Information**

Application Primary Accession	WB, IF, ICC <u>Q9UJX2</u> Det Human Maure
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CDC23; ANAPC8; APC8; Cell division cycle 23; CUT23; Cyclosome subunit 8;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	68834

## **Additional Information**

Dilution Purification Immunogen Description	WB 1:500~1:2000 ICC/IF 1:50~1:100 Affinity-chromatography A synthesized peptide derived from human Cdc23/APC8 Anaphase-promoting complex subunit 8 (APC8, CDC23) is a component of the tetratricopeptide repeat (TPR) APC/C sub-complex that also includes APC3 (CDC27) and APC6 (CDC16). APC8 protein associates with APC3 and APC6 to facilitate recruitment of the APC/C coactivation subunits CDC20 and Cdh1/FZR1. Research studies suggest that APC8 protein is overexpressed in papillary thyroid cancer and acts as an important regulator of cell cycle
Storage Condition and Buffer	progression and cell growth.

## **Protein Information**

Name	CDC23
Synonyms	ANAPC8
Function	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle (PubMed: <u>18485873</u> ). The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed: <u>18485873</u> ). The APC/C complex catalyzes assembly of branched 'Lys-11'-/'Lys-48'-linked branched ubiquitin chains on target proteins (PubMed: <u>29033132</u> ).



Western blot analysis of Cdc23/APC8 expression in (1)HepG2 cell lysate; (2)Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.