

## PAX5 Antibody

Rabbit mAb Catalog # AP90969

## **Product Information**

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IF, FC, ICC, IHF <u>Q02548</u> Human, Mouse Monoclonal B cell specific activator protein; B-cell lineage specific activator; BSAP; paired box 5; paired box homeotic gene 5; Paired box protein Pax-5; PAX5; transcription factor PAX 5;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	42149

## **Additional Information**

Dilution Purification Immunogen	WB 1:1000~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:30 Affinity-chromatography A synthesized peptide derived from human PAX5
Description	AX5 May play an important role in B-cell differentiation as well as neural development and spermatogenesis. Involved in the regulation of the CD19 gene, a B-lymphoid-specific target gene. Interacts with DAXX. Binds DNA as a monomer. Binds TLE4.
Storage Condition and Buffer	

## **Protein Information**

Name	PAX5
Function	Transcription factor that plays an essential role in commitment of lymphoid progenitors to the B-lymphocyte lineage (PubMed: <u>10811620</u> , PubMed: <u>27181361</u> ). Fulfills a dual role by repressing B-lineage inappropriate genes and simultaneously activating B-lineage- specific genes (PubMed: <u>10811620</u> , PubMed: <u>27181361</u> ). In turn, regulates cell adhesion and migration, induces V(H)-to-D(H)J(H) recombination, facilitates pre-B-cell receptor signaling and promotes development to the mature B-cell stage (PubMed: <u>32612238</u> ). Repression of the cohesin- release factor WAPL causes global changes of the chromosomal architecture in pro-B cells to facilitate the generation of a diverse antibody repertoire (PubMed: <u>32612238</u> ).
Cellular Location	Nucleus.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.