

POU2AF1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9096a

Product Information

Application	FC, IHC-P, WB, E
Primary Accession	Q16633
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB22378
Calculated MW	27436
Antigen Region	1-27

Additional Information

Gene ID	5450
Other Names	POU domain class 2-associating factor 1, B-cell-specific coactivator OBF-1, BOB-1, OCA-B, OCT-binding factor 1, POU2AF1, OBF1
Target/Specificity	This POU2AF1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-27 amino acids from the N-terminal region of human POU2AF1.
Dilution	FC~~1:10~50 IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	POU2AF1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	POU2AF1 (HGNC:9211)
Function	Transcriptional coactivator that specifically associates with either POU2F1/OCT1 or POU2F2/OCT2 (PubMed: 7859290). It boosts the POU2F1/OCT1 mediated promoter activity and to a lesser extent, that of

POU2F2/OCT2 (PubMed:[7779176](#)). It recognizes the POU domains of POU2F1/OCT1 and POU2F2/OCT2 (PubMed:[7779176](#)). It is essential for the response of B-cells to antigens and required for the formation of germinal centers (PubMed:[7623806](#), PubMed:[7859290](#)). Regulates IL6 expression in B cells as POU2F2/OCT2 coactivator (By similarity).

Cellular Location

Nucleus.

Tissue Location

B-cell specific (PubMed:[7779176](#), PubMed:[7859290](#)). Detected in mainly in spleen, but also in thymus, peripheral blood leukocyte and small intestine (PubMed:[7779176](#), PubMed:[7859290](#))


Background

POU2AF1 is transcriptional coactivator that specifically associates with either OCT1 or OCT2. It boosts the OCT1 mediated promoter activity and to a lesser extent, that of OCT2. It has no intrinsic DNA-binding activity. It recognizes the POU domains of OCT1 and OCT2. It is essential for the response of B-cells to antigens and required for the formation of germinal centers.


References

Pittman,A.M., et.al., Hum. Mol. Genet. 17 (23), 3720-3727 (2008)
Gashaw,I., et.al., Mol. Hum. Reprod. 13 (10), 721-727 (2007)

Images




Anti-POU2AF1 Antibody (N-term) at 1:2000 dilution + Ramos whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



POU2AF1 Antibody (N-term) (Cat. #AP9096a) flow cytometry analysis of Ramos cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

POU2AF1 Antibody (N-term) (Cat. #AP9096a)immunohistochemistry analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the



use of POU2AF1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.