

# KIF11 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9097A

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P52732</a>
<b>Reactivity</b>	Human, Hamster
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB22568
<b>Calculated MW</b>	119159
<b>Antigen Region</b>	103-130

## Additional Information

---

<b>Gene ID</b>	3832
<b>Other Names</b>	Kinesin-like protein KIF11, Kinesin-like protein 1, Kinesin-like spindle protein HKSP, Kinesin-related motor protein Eg5, Thyroid receptor-interacting protein 5, TR-interacting protein 5, TRIP-5, KIF11, EG5, KNSL1, TRIP5
<b>Target/Specificity</b>	This KIF11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 103-130 amino acids from the N-terminal region of human KIF11.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	KIF11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	KIF11
<b>Synonyms</b>	EG5, KNSL1, TRIP5

<b>Function</b>	Motor protein required for establishing a bipolar spindle and thus contributing to chromosome congression during mitosis (PubMed: <a href="#">19001501</a> , PubMed: <a href="#">37728657</a> ). Required in non-mitotic cells for transport of secretory proteins from the Golgi complex to the cell surface (PubMed: <a href="#">23857769</a> ).
<b>Cellular Location</b>	Cytoplasm. Cytoplasm, cytoskeleton, spindle pole


## Background

KIF11 belongs to the kinesin-like protein family. Members of this protein family are known to be involved in various kinds of spindle dynamics. The function of this gene product includes chromosome positioning, centrosome separation and establishing a bipolar spindle during cell mitosis.


## References

Tcherniuk,S., et.al., Biochem. Pharmacol. 79 (6), 864-872 (2010)  
 Parke,C.L., et.al., J. Biol. Chem. 285 (8), 5859-5867 (2010)

## Images





Anti-KIF11 Antibody (N-term) at 1:1000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 119 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of KIF11 Antibody (N-term) (Cat. #AP9097a) in CHO cell line lysates (35ug/lane). KIF11 (arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human lung carcinoma reacted with KIF11 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



KIF11 Antibody (N-term) (Cat.#AP9097a) flow cytometry analysis of Hela cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.