10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SUZ12 Antibody

Rabbit mAb
Catalog # AP90989

Product Information

Application	WB, IF, ICC, IP
Primary Accession	Q15022
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CHET9; JJAZ1; KIAA0160; SUZ12;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	83055

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SUZ12
Description	Polycomb group (PcG) protein. Component of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene. The PRC2/EED-EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SUZ12
Synonyms	CHET9, JJAZ1, KIAA0160
Function	Polycomb group (PcG) protein. Component of the PRC2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene (PubMed:15225548, PubMed:15231737, PubMed:15385962, PubMed:16618801, PubMed:17344414, PubMed:18285464, PubMed:28229514, PubMed:29499137, PubMed:31959557). The PRC2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems (PubMed:12351676, PubMed:12435631, PubMed:15099518, PubMed:15225548, PubMed:15385962, PubMed:15684044, PubMed:16431907, PubMed:18086877, PubMed:18285464). Genes repressed by the PRC2 complex include HOXC8, HOXA9, MYT1 and CDKN2A (PubMed:15231737, PubMed:16618801,

PubMed:17200670, PubMed:31959557).

Cellular Location	Nucleus Note=Localizes to chromatin as part of the PRC2 complex
Tissue Location	Overexpressed in breast and colon cancer.

Images



Western blot analysis of SUZ12 expression in (1) HeLa cell lysate; (2) SW480 cell lysate.