

# Fas Antibody

Rabbit mAb Catalog # AP91031

## **Product Information**

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	<u>P25445</u>
Reactivity	Human
Clonality	Monoclonal
Other Names	FAS; ALPS1A; APO-1; APT1; CD95; FAS1; FASTM; TNFRSF6;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	37732

### **Additional Information**

Dilution Purification Immunogen	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50 Affinity-chromatography A synthesized peptide derived from human Fas
Description	Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis.
Storage Condition and Buffer	

#### **Protein Information**

Name	FAS
Synonyms	APT1, FAS1, TNFRSF6
Function	Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase CASP8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs CASP8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen- stimulated suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Membrane raft [Isoform 3]: Secreted. [Isoform 5]: Secreted.
Tissue Location	Isoform 1 and isoform 6 are expressed at equal levels in resting peripheral

blood mononuclear cells. After activation there is an increase in isoform 1 and decrease in the levels of isoform 6.

# Images



Western blot analysis of Fas expression in Ramos cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.