

HMGR Antibody

Rabbit mAb

Catalog # AP91071

Product Information

Application	WB, IP
Primary Accession	P04035
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	3H3M; HMDH; HMG CoA reductase; HMG CoAR;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	97476


Additional Information

Dilution	WB 1:500~1:2000 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HMGR
Description	Transmembrane glycoprotein that is the rate-limiting enzyme in cholesterol biosynthesis as well as in the biosynthesis of nonsterol isoprenoids that are essential for normal cell function including ubiquinone and geranylgeranyl proteins.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	HMGR (HGNC:5006)
Function	Catalyzes the conversion of (3S)-hydroxy-3-methylglutaryl-CoA (HMG-CoA) to mevalonic acid, the rate-limiting step in the synthesis of cholesterol and other isoprenoids, thus plays a critical role in cellular cholesterol homeostasis (PubMed: 21357570 , PubMed: 2991281 , PubMed: 36745799 , PubMed: 6995544). HMGR is the main target of statins, a class of cholesterol-lowering drugs (PubMed: 11349148 , PubMed: 18540668 , PubMed: 36745799).
Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P00347}. Peroxisome membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P00347}
Tissue Location	[Isoform 1]: Ubiquitously expressed with the highest levels in the cerebellum, fetal brain, testis, skin and adrenal gland. [Isoform 3]: Low abundance except in skin, esophagus, and uterine cervix.

Images



Western blot analysis of HMGCR expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.