

# Ctip2 Antibody

Rabbit mAb Catalog # AP91081

## **Product Information**

Application Primary Accession Reactivity Clonality Other Names	WB <u>Q9C0K0</u> Human Monoclonal ATL1 alpha; ATL1 beta; ATL1 gamma; ATL1 delta; Bcl11b; BCL11B/TRDC fusion; CTIP2; hRIT1 alpha; hRit1; Rit1;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	95519


### **Additional Information**

Dilution Purification Immunogen	WB 1:500~1:2000 Affinity-chromatography A synthesized peptide derived from human Ctip2
Description	Tumor-suppressor protein involved in T-cell lymphomas. May function on the P53-signaling pathway. May be a key regulator of both differentiation and survival during thymocyte development. Repress transcription through direct, TFCOUP2-independent binding to a GC-rich response element.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

#### **Protein Information**

Name	BCL11B
Synonyms	CTIP2, RIT1
Function	Key regulator of both differentiation and survival of T- lymphocytes during thymocyte development in mammals. Essential in controlling the responsiveness of hematopoietic stem cells to chemotactic signals by modulating the expression of the receptors CCR7 and CCR9, which direct the movement of progenitor cells from the bone marrow to the thymus (PubMed: <u>27959755</u> ). Is a regulator of IL2 promoter and enhances IL2 expression in activated CD4(+) T-lymphocytes (PubMed: <u>16809611</u> ). Tumor-suppressor that represses transcription through direct, TFCOUP2-independent binding to a GC-rich response element (By similarity). May also function in the P53-signaling pathway (By similarity).
Cellular Location	Nucleus.

## Images



Western blot analysis of Ctip2 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.