

PGP9.5 Antibody

Rabbit mAb Catalog # AP91153

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IF, FC, ICC, IP, IHF <u>P09936</u> Rat, Human, Mouse Monoclonal HEL 117; NDGOA; Neuron cytoplasmic protein 9.5; PARK5; PGP9.5; Protein gene product 9.5; UCHL1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	24824

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:50 Affinity-chromatography A synthesized peptide derived from human PGP9.5 Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. Also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer may have ATP-independent ubiquitin ligase
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name

UCHL1

FunctionDeubiquitinase that plays a role in the regulation of several processes such
as maintenance of synaptic function, cardiac function, inflammatory response
or osteoclastogenesis (PubMed:22212137, PubMed:23359680). Abrogates the
ubiquitination of multiple proteins including WWTR1/TAZ, EGFR, HIF1A and
beta-site amyloid precursor protein cleaving enzyme 1/BACE1
(PubMed:22212137, PubMed:25615526). In addition, recognizes and
hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin to maintain a
stable pool of monoubiquitin that is a key requirement for the
ubiquitin-proteasome and the autophagy- lysosome pathways
(PubMed:12408865, PubMed:8639624, PubMed:9774100). Regulates amyloid
precursor protein/APP processing by promoting BACE1 degradation resulting
in decreased amyloid beta production (PubMed:22212137). Plays a role in the

	immune response by regulating the ability of MHC I molecules to reach cross-presentation compartments competent for generating Ag-MHC I complexes (By similarity). Mediates the 'Lys-48'-linked deubiquitination of the transcriptional coactivator WWTR1/TAZ leading to its stabilization and inhibition of osteoclastogenesis (By similarity). Deubiquitinates and stabilizes epidermal growth factor receptor EGFR to prevent its degradation and to activate its downstream mediators (By similarity). Modulates oxidative activity in skeletal muscle by regulating key mitochondrial oxidative proteins (By similarity). Enhances the activity of hypoxia-inducible factor 1-alpha/HIF1A by abrogateing its VHL E3 ligase-mediated ubiquitination and consequently inhibiting its degradation (PubMed:25615526).
Cellular Location	Cytoplasm. Endoplasmic reticulum membrane; Lipid- anchor. Note=About 30% of total UCHL1 is associated with membranes in brain. Localizes near and/or within mitochondria to potentially interact with mitochondrial proteins {ECO:0000250 UniProtKB:Q9R0P9}
Tissue Location	Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.

Images



Western blot analysis of PGP9.5 expression in 293T cell lysate.

Image not found : 202311/AP91153-IHC.jpg	Immunohistochemical analysis of paraffin-embedded human prostate cancer, using PGP9.5 Antibody.
Image not found : 202311/AP91153-IF.jpg	Immunofluorescent analysis of U87-MG cells, using PGP9.5 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.