

Aromatase Antibody

Rabbit mAb Catalog # AP91162

Product Information

| Application Primary Accession Reactivity Clonality Other Names | WB, IP <u>P11511</u> Rat, Human Monoclonal Aromatase; CYPXIX; Cytochrome P-450AROM; Cytochrome P450 19A1; Estrogen synthase; CYP19A1; ARO1; CYAR; CYP19; |
|--|---|
| lsotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 57883 |

Additional Information

| Dilution | WB 1:500~1:2000 IP 1:50 |
|------------------------------|---|
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human Aromatase |
| Description | Catalyzes the formation of aromatic C18 estrogens from C19 androgens. |
| Storage Condition and Buffer | , |

Protein Information

| Name | CYP19A1 {ECO:0000303 PubMed:24705274, ECO:0000312 HGNC:HGNC:2594} |
|----------|--|
| Function | A cytochrome P450 monooxygenase that catalyzes the conversion of C19 androgens, androst-4-ene-3,17-dione (androstenedione) and testosterone to the C18 estrogens, estrone and estradiol, respectively (PubMed: <u>27702664</u> , PubMed: <u>2848247</u>). Catalyzes three successive oxidations of C19 androgens: two conventional oxidations at C19 yielding 19-hydroxy and 19-oxo/19-aldehyde derivatives, followed by a third oxidative aromatization step that involves C1-beta hydrogen abstraction combined with cleavage of the C10-C19 bond to yield a phenolic A ring and formic acid (PubMed: <u>20385561</u>). Alternatively, the third oxidative reaction yields a 19-norsteroid and formic acid. Converts dihydrotestosterone to delta1,10-dehydro 19- nordihydrotestosterone and may play a role in homeostasis of this potent androgen (PubMed: <u>22773874</u>). Also displays 2-hydroxylase activity toward estrone (PubMed: <u>22773874</u>). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein |

reductase) (PubMed:20385561, PubMed:22773874).Cellular LocationEndoplasmic reticulum membrane; Multi-pass membrane protein. Microsome
membrane; Multi- pass membrane proteinTissue LocationWidely expressed, including in adult and fetal brain, placenta, skin fibroblasts,
adipose tissue and gonads

Images



Western blot analysis of Aromatase expression in rat brain lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.