

# Rhodopsin Antibody

Rabbit mAb

Catalog # AP91190

## Product Information

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">P08100</a>
<b>Reactivity</b>	Rat, Human
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	CSNBAD1; OPN2; opsd; Opsin 2 rod pigment; Opsin2; Retinitis pigmentosa 4 autosomal dominant; RHO; Rhodopsin; RP4;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	38893

## Additional Information

<b>Dilution</b>	WB 1:500~1:2000 IHC 1:50~1:200
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human Rhodopsin
<b>Description</b>	Photoreceptor required for image-forming vision at low light intensity. Required for photoreceptor cell viability after birth. Light-induced isomerization of 11-cis to all-trans retinal triggers a conformational change leading to G-protein activation and release of all-trans retinal.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	RHO
<b>Synonyms</b>	OPN2
<b>Function</b>	Photoreceptor required for image-forming vision at low light intensity (PubMed: <a href="#">7846071</a> , PubMed: <a href="#">8107847</a> ). Required for photoreceptor cell viability after birth (PubMed: <a href="#">12566452</a> , PubMed: <a href="#">2215617</a> ). Light- induced isomerization of the chromophore 11-cis-retinal to all-trans- retinal triggers a conformational change that activates signaling via G-proteins (PubMed: <a href="#">26200343</a> , PubMed: <a href="#">28524165</a> , PubMed: <a href="#">28753425</a> , PubMed: <a href="#">8107847</a> ). Subsequent receptor phosphorylation mediates displacement of the bound G-protein alpha subunit by the arrestin SAG and terminates signaling (PubMed: <a href="#">26200343</a> , PubMed: <a href="#">28524165</a> ).
<b>Cellular Location</b>	Membrane; Multi-pass membrane protein. Cell projection, cilium, photoreceptor outer segment. Note=Synthesized in the inner segment (IS) of


rod photoreceptor cells before vectorial transport to disk membranes in the rod outer segment (OS) photosensory cilia

**Tissue Location**

Rod shaped photoreceptor cells which mediate vision in dim light

**Images**

---



Western blot analysis of Rhodopsin expression in rat eyeball lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.