

PRDX6 Antibody

Rabbit mAb Catalog # AP91214

Product Information

Application WB, IF, ICC, IP

Primary Accession <u>P30041</u>

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names Peroxiredoxin-6; Antioxidant protein 2; Liver 2D page spot 40; NSGPx; PRDX6;

AOP2; KIAA;

IsotypeRabbit IgGHostRabbitCalculated MW25035

Additional Information

Dilution WB 1:500~1:1000 ICC/IF 1:50~1:200 IP 1:20

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human PRDX6

Description Involved in redox regulation of the cell. Can reduce H(2)O(2) and short chain

organic, fatty acid, and phospholipid hydroperoxides. May play a role in the regulation of phospholipid turnover as well as in protection against oxidative

injury.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name PRDX6

Synonyms AOP2, KIAA0106

Function Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide

and organic hydroperoxides to water and alcohols, respectively

(PubMed: 10893423, PubMed: 9497358). Can reduce H(2)O(2) and short chain organic, fatty acid, and phospholipid hydroperoxides (PubMed: 10893423). Also has phospholipase activity, can therefore either reduce the oxidized sn-2 fatty acyl group of phospholipids (peroxidase activity) or hydrolyze the sn-2 ester bond of phospholipids (phospholipase activity) (PubMed: 10893423,

PubMed: 26830860). These activities are dependent on binding to phospholipids at acidic pH and to oxidized phospholipds at cytosolic pH (PubMed: 10893423). Plays a role in cell protection against oxidative stress by detoxifying peroxides and in phospholipid homeostasis (PubMed: 10893423).


Exhibits acyl-CoA-dependent lysophospholipid acyltransferase which

mediates the conversion of lysophosphatidylcholine (1-acyl-sn-glycero-3-phosphocholine or LPC) into phosphatidylcholine (1,2-diacyl-sn-glycero-3-phosphocholine or PC) (PubMed:26830860). Shows a clear preference for LPC as the lysophospholipid and for palmitoyl CoA as the fatty acyl substrate (PubMed:26830860).

Cellular Location

Cytoplasm. Lysosome {ECO:0000250|UniProtKB:O35244}. Note=Also found in lung secretory organelles (lamellar bodies). {ECO:0000250|UniProtKB:O35244}

Images



Western blot analysis of PRDX6 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.