

# CD40L Antibody

Rabbit mAb

Catalog # AP91237

## Product Information

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P29965</a>
<b>Reactivity</b>	Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	CD40 ligand; CD40-L; T-cell antigen Gp39; TNF-related activation protein; TRAP; CD154; CD40 ligand, membrane form; CD40 ligand, soluble form; CD40LG; CD40L; TNFSF5; TRAP;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	29274

## Additional Information

<b>Dilution</b>	WB 1:500~1:2000
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human CD40L
<b>Description</b>	Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL-4. Involved in immunoglobulin class switching.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	CD40LG
<b>Synonyms</b>	CD40L, TNFSF5, TRAP
<b>Function</b>	Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed: <a href="#">1280226</a> , PubMed: <a href="#">31331973</a> ). Costimulates T-cell proliferation and cytokine production (PubMed: <a href="#">8617933</a> ). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed: <a href="#">8617933</a> ). Induces the activation of NF-kappa-B (PubMed: <a href="#">15067037</a> , PubMed: <a href="#">31331973</a> ). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed: <a href="#">15067037</a> ). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed: <a href="#">15067037</a> ). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).

**Cellular Location**


Cell membrane; Single-pass type II membrane protein. Cell surface

**Tissue Location**

Specifically expressed on activated CD4+ T- lymphocytes

**Images**

---



Western blot analysis of CD40L expression in MOLT-4 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.