

CD73 Antibody

Rabbit mAb

Catalog # AP91243

Product Information

Application	WB, IHC
Primary Accession	P21589
Reactivity	Human
Clonality	Monoclonal
Other Names	5' nucleotidase (CD73); 5' nucleotidase precursor; 5' nucleotidase; ecto; 5NTD; CD73; CD73 antigen; E5NT; Ecto 5' nucleotidase; Ecto-5'-nucleotidase; NT5E; NTE;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	63368


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD73
Description	Hydrolyzes extracellular nucleotides into membrane permeable nucleosides.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	NT5E
Synonyms	NT5, NTE
Function	Catalyzes the hydrolysis of nucleotide monophosphates, releasing inorganic phosphate and the corresponding nucleoside, with AMP being the preferred substrate (PubMed: 21933152 , PubMed: 22997138 , PubMed: 23142347 , PubMed: 24887587 , PubMed: 34403084). Shows a preference for ribonucleotide monophosphates over their equivalent deoxyribose forms (PubMed: 34403084). Other substrates include IMP, UMP, GMP, CMP, dAMP, dCMP, dTMP, NAD and NMN (PubMed: 21933152 , PubMed: 22997138 , PubMed: 23142347 , PubMed: 24887587 , PubMed: 34403084).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor

Images



Western blot analysis of CD73 expression in A375 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.